

On the existence of stable models of non-stratified logic programs

Stefania COSTANTINI

Dip. di Informatica, Università di L'Aquila
via Vetoio Loc. Coppito, L'Aquila, I-67010 Italy
(e-mail: stefcost@di.univaq.it)

submitted 11 June 2000; revised 6 November 2003, 15 July 2004; accepted 20 December 2004

Abstract

In this paper we analyze the relationship between cyclic definitions and consistency in Gelfond-Lifschitz's answer sets semantics (originally defined as 'stable model semantics'). This paper introduces a fundamental result, which is relevant for Answer Set programming, and planning. For the first time since the definition of the stable model semantics, the class of logic programs for which a stable model exists is given a syntactic characterization. This condition may have a practical importance both for defining new algorithms for checking consistency and computing answer sets, and for improving the existing systems. The approach of this paper is to introduce a new *canonical form* (to which any logic program can be reduced to), to focus the attention on cyclic dependencies. The technical result is then given in terms of programs in canonical form (canonical programs), without loss of generality: the stable models of any general logic program coincide (up to the language) to those of the corresponding canonical program. The result is based on identifying the cycles contained in the program, showing that stable models of the overall program are composed of stable models of suitable sub-programs, corresponding to the cycles, and on defining the *Cycle Graph*. Each vertex of this graph corresponds to one cycle, and each edge corresponds to one *handle*, which is a literal containing an atom that, occurring in both cycles, actually determines a connection between them. In fact, the truth value of the handle in the cycle where it appears as the head of a rule, influences the truth value of the atoms of the cycle(s) where it occurs in the body. We can therefore introduce the concept of a *handle path*, connecting different cycles. Cycles can be even, if they consist of an even number of rules, or vice versa they can be odd. Problems for consistency, as it is well-known, originate in the odd cycles. If for every odd cycle we can find a handle path with certain properties, then the existence of stable model is guaranteed. We will show that based on this results new classes of consistent programs can be defined, and that cycles and cycle graphs can be generalized to components and *component graphs*.

1 Introduction

In this paper we analyze the relationship between cyclic definitions and consistency in Gelfond-Lifschitz's answer sets semantics. As it is well-known, under the answer set semantics a theory may have no *answer sets*, since the corresponding general logic program may have no *stable models* (Gelfond and Lifschitz 1988) (Gelfond and Lifschitz 1991).

We introduce a fundamental result, which is relevant for Answer Set Programming (Marek and Truszczyński 1999), (Niemelä 1999) planning (Lifschitz 1999) and diagnosis (Balduccini and Gelfond 2003). For the first time, the class of logic programs for which a stable model exists is given a syntactic characterization (the result extends naturally to answer sets semantics) by providing a necessary and sufficient condition.

While checking for the existence of stable models is as hard a computational problem (in fact, NP-complete) as planning under certain assumptions (see (Liberatore 1999)), consistency checking is a good conceptual tool when derivations are based on consistency arguments. This is the case for instance for all formalizations that treat goals as constraints over models of the program. Then, being able to check for the existence of stable models syntactically for every answer set program can be of help for the logic programming encodings of planning and diagnosis (like, e.g., those of (Erdem and Lifschitz 1999), (Faber et al. 1999), (Balduccini et al. 2000), (Dimopoulos et al. 1997) and (Balduccini and Gelfond 2003)), and for Answer Set Programming in general.

The approach of this paper is to introduce a new *canonical form* to which any logic program can be reduced. The technical result is then given in terms of programs in canonical form (canonical programs), without loss of generality. Canonical programs focus the attention on cyclic dependencies. Rules are kept short, so as to make the syntactic analysis of the program easier. The stable models of any general logic program coincide (up to the language) to those of the corresponding canonical program.

A detailed analysis of the steps involved in reducing programs to their canonical form has been performed in (Costantini and Proveti 2004) and, as intuition suggests, this transformation is tractable. Nevertheless, all definitions and results presented in this paper might be rephrased for general programs without conceptual problems, just at the expense of a lot of additional details. This means that reduction to canonical form is not strictly required neither for the theory, nor for an implementation.

The main result of this paper is a necessary and sufficient syntactic condition for the existence of stable models. On the one hand, this condition is of theoretical interest, as it is the first one ever defined since the introduction of the stable model semantics in (Gelfond and Lifschitz 1988). On the other hand, it may have a practical importance both for defining new algorithms for checking consistency and computing answer sets, and for improving the existing systems (Solvers 2004).

The result is based on identifying the cycles contained in the program, on showing that stable models of the overall program are composed of stable models of suitable sub-programs, corresponding to the cycles, and on representing the program by means of its *Cycle Graph*. Each vertex of this graph corresponds to one cycle, and each edge corresponds to one *handle*, which is a literal containing an atom that, occurring in both cycles,

actually determines a connection between them. In fact, the truth value of the handle in the cycle where it appears as the head of a rule influences the truth value of the atoms of the cycle(s) where it occurs in the body. We can therefore introduce the concept of a *handle path*, connecting different cycles. Cycles can be even, if they consist of an even number of rules, or vice versa they can be odd. Problems for consistency, as it is well-known, originate in the odd cycles. If and only if for every odd cycle we can find a subgraph with certain properties, then the existence of stable models is guaranteed.

The necessary and sufficient condition that we introduce is syntactic in the sense that it does not refer either to models or derivations. Checking this condition requires neither finding the stable models nor applying the rules of the program. It just requires to look at the program (represented by the Cycle Graph) and at the rules composing the cycles. The condition can however be exploited, so as to obtain: (i) new algorithms for finding the stable models, which are at least of theoretical interest; (ii) a new method for consistency checking divided into two steps: a first step related to the coarse structure of the program, that can be easily checked on the Cycle Graph so as to rule out a lot of inconsistent programs, thus leaving only the potentially consistent ones to be verified in a second step, that can be performed according to the approach presented here, or in any other way.

We will argue that the approach can also be useful for defining classes of programs that are consistent by construction, and as a first step toward a component-based methodology for the construction and analysis of answer set programs. This by means of a further generalization of Cycle Graphs to *Component Graphs*, where vertices are *components* consisting of bunches of cycles, and edges connect different components. We will argue that, in this framework, components can even be understood as independent agents.

It is useful to notice that in Answer Set Programming graph representations have been widely used for studying and characterizing properties of answer set programs, first of all consistency, and for computing the answer sets. Among the most important approaches we may mention the Rule Graph (Dimopoulos and Torres 1996) and its extensions (Linke 2001) (Linke 2003b) (Konczak et al. 2003a), and the Extended Dependency Graph (Brignoli et al. 1999), that we have considered and compared (Costantini 2001), (Costantini et al. 2002). Enhanced classes of graphs have been recently introduced in order to cope with extensions to the basic paradigm such as for instance preferences (Konczak et al. 2003b) or nested logic programs (Linke 2003a). However, the Cycle Graph proposed in this paper is different from all the above-mentioned approaches, since its vertices are not atoms or rules, but significant subprograms (namely cycles), and the edges are connections between these subprograms.

2 Preliminary Definitions

We consider the standard definition of a (propositional) general logic program and of well-founded (Van Gelder et al. 1990) and stable model (Gelfond and Lifschitz 1988) and answer set semantics (Gelfond and Lifschitz 1991). Whenever we mention consistency (or stability) conditions, we refer to the conditions introduced in (Gelfond and Lifschitz 1988). This section summarizes some basic definitions, and is intended for readers who are unfamiliar with the above-mentioned topics.

Assume a language of constants and predicate constants. Assume also that terms and atoms are built as in the corresponding first-order language. Unlike classical logic and standard logic programming, no function symbols are allowed. A rule is an expression of the form:

$$\rho : \lambda_0 \leftarrow \lambda_1, \dots, \lambda_m, \text{not } \lambda_{m+1}, \dots, \text{not } \lambda_n \quad (1)$$

where $\lambda_0, \dots, \lambda_n$ are atoms and *not* is a logical connective called *negation as failure*. The λ_i 's are called *positive literals*, and the *not* λ_j 's *negative literals*. For every rule let us define $\text{head}(\rho) = \lambda_0$ (also called the *conclusion* of the rule), $\text{pos}(\rho) = \lambda_1, \dots, \lambda_m$, $\text{neg}(\rho) = \lambda_{m+1}, \dots, \lambda_n$ and $\text{body}(\rho) = \text{pos}(\rho) \cup \text{neg}(\rho)$ (also called the *conditions* of the rule). If $\text{body}(\rho) = \emptyset$ we refer to ρ as a *unit rule* (w.r.t. non-unit rules), or a *fact*. We will say that $\text{head}(\rho)$ *depends on*, or *is defined in terms of*, the literals in $\text{body}(\rho)$.

A general logic program Π (or simply “logic program”) is defined as a collection of rules. In the rest of this paper, we rely on the assumption that the order of literals in the body of rules is irrelevant. Rules with variables are taken as shorthand for the sets of all their ground instantiations and the set of all ground atoms in the language of a program Π will be denoted by $\mathbb{B}_\Pi$.

2.1 Semantics

For the sake of simplicity, we give here the definition of *stable model* instead of that of answer set, which is an extension given for programs that contain the explicit negation operator $\neg$. In fact, this is not going to make a difference in the context of this work, and we will often interchange the terms “stable models” and “answer sets”. Intuitively, a stable model is a possible view of the world that is *compatible* with the rules of the program. Rules are therefore seen as constraints on these views of the world.

Let us start by defining stable models of the subclass of positive programs, i.e. those where, for every rule ρ , $\text{neg}(\rho) = \emptyset$.

Definition 1

(Stable Models of positive logic programs)

The *stable model* $a(\Pi)$ of a positive program Π is the smallest subset of $\mathbb{B}_\Pi$ such that for any rule (1) in Π :

$$\lambda_1, \dots, \lambda_m \in a(\Pi) \Rightarrow \lambda_0 \in a(\Pi) \quad (2)$$

Positive programs have a unique stable model, which coincides with its minimal model, that can also be obtained applying other semantics; then, positive programs are unambiguous. The stable model of a positive program can be obtained as the fixed point of the *immediate consequence operator*

$$T_\Pi(I) = \{\lambda : \exists \rho \in \Pi \text{ s.t. } \lambda = \text{head}(\rho) \wedge \text{pos}(\rho) \subseteq I\}$$

The iterated application of T_Π from $\emptyset$ on (i.e., $T_\Pi(\emptyset), T_\Pi^2(\emptyset), \dots$) is guaranteed to have a fixed point, which corresponds to the unique stable model (answer set) of Π .

A set of atoms $\mathcal{S}$ is a stable model of an (arbitrary) program if it is a minimal model and every atom $\alpha \in \mathcal{S}$ is supported by some rule of the program. With respect of negation, if we assume $\mathcal{S}$ to be a stable model: (i) no atom can belong to $\mathcal{S}$, which is derived by means of a rule with a condition *not* α where α is true in $\mathcal{S}$, i.e. $\alpha \in \mathcal{S}$; (ii) all literals *not* β in the body of rules where β is false in $\mathcal{S}$ are, of course, true in $\mathcal{S}$. Consequently, in order to check whether $\mathcal{S}$ actually is a stable model, all negations can be deleted according to these criteria, in order to apply the above formulation for positive programs.

Definition 2

(Stable Model of arbitrary logic programs)

Let Π be a logic program. For any set $\mathcal{S}$ of atoms, let $\Pi^\mathcal{S}$ be a program obtained from Π by deleting

- (i) each rule that has a formula “*not* λ ” in its body with $\lambda \in \mathcal{S}$;
- (ii) all formulae of the form “*not* λ ” in the bodies of the remaining rules.

Since $\Pi^\mathcal{S}$ does not contain *not*, its stable model is already defined. If this stable model coincides with $\mathcal{S}$, then we say that $\mathcal{S}$ is a *stable model* of Π . Precisely, a stable model of Π is characterized by the equation:

$$\mathcal{S} = a(\Pi^\mathcal{S}). \quad (3)$$

The Γ operator, introduced by Gelfond and Lifschitz in (Gelfond and Lifschitz 1988), is defined as $\Gamma(\Pi, \mathcal{S}) = a(\Pi^\mathcal{S})$. When Π is fixed, we may drop the first parameter and refer to Γ as a function from the powerset of $\mathbb{B}_\Pi$ to itself. In practice however, stable models are not computed by applying Γ to all subsets of $\mathbb{B}_\Pi$. Answer set solvers (Solvers 2004) in fact apply more effective and smarter algorithms.

Stable models are minimal models of Π in the classical sense, but the converse does not hold. Then, a program may have no stable models. In general a program has several stable models, and programs with a unique stable model are called *categorical*. In this paper, consistency means existence of stable models (or, equivalently, of answer sets). Conventionally, “an atom being true” means “an atom being in a stable model”. Whenever we consider a set of atoms $\mathcal{I}$, we implicitly mean $\mathcal{I} \subseteq \mathbb{B}_\Pi$. We say that a literal α (respectively *not* α) is true w.r.t. $\mathcal{I}$ if $\alpha \in \mathcal{I}$ (respectively $\mathcal{I}$ if $\alpha \notin \mathcal{I}$).

2.2 The well-founded semantics

The well-founded semantics of (Van Gelder et al. 1990) assigns to a logic program Π a unique, three-valued model, called well-founded model and denoted by $WFS(\Pi) = \langle W^+, W^- \rangle$, where W^+ and W^- are disjoint. Intuitively, W^+ is the set of atoms deemed true, W^- is the set of atoms deemed false, while atoms belonging to neither set are deemed *undefined*.

The reduction of a program to its *canonical form* that we discuss later is based on a preliminary simplification of the program w.r.t. the well-founded semantics. The result of this is a compact version (or *reduct*) of the program which is *WF-irreducible*, where

Definition 3

A program Π is *WF-irreducible* if and only if $WFS(\Pi) = \langle \emptyset, \emptyset \rangle$.

For general logic programs, atoms with truth value “undefined” under the well-founded semantics are exactly the atoms which are of interest for finding the stable models. This is a consequence of the fact that all stable models of a program *extend* its well-founded model, i.e., every literal which is true (resp. false) in the well-founded model is also true (resp. false) in all stable models. The stable models of the original program can be easily obtained from the stable models of the WF-irreducible reduct (Costantini 1995), and vice versa, if the reduct has no stable models the same holds for the original program.

For instance, for program

$$p \leftarrow \text{not } p, \text{not } q$$

with well-founded model $\langle \emptyset; \{q\} \rangle$ where atom p has truth value “undefined”, we get the simplified WF-irreducible reduct $p \leftarrow \text{not } p$ by getting rid of a literal which is true under the well-founded semantics, and thus is true in all stable models (if any exists). The reduct has no stable models, like the original program.

3 Cycles and Handles

The results on consistency checking that we will present later are based on identifying the negative cycles contained in the program, on showing that stable models of the overall program are composed of stable models of suitable sub-programs, corresponding to the negative cycles, and on representing the program by means of its *Cycle Graph*. In this section we define when a set of rules constitutes a negative cycle (or simply “cycle”), which kinds of cycles we may have and how cycles can be understood to be connected to each other.

Definition 4

A set of rules C is called a *negative cycle*, or for short a *cycle*, if it has the following form:

$$\begin{aligned} \lambda_1 &\leftarrow \text{not } \lambda_2, \Delta_1 \\ \lambda_2 &\leftarrow \text{not } \lambda_3, \Delta_2 \\ &\dots \\ \lambda_n &\leftarrow \text{not } \lambda_1, \Delta_n \end{aligned}$$

where $\lambda_1, \dots, \lambda_n$ are distinct atoms. Each Δ_i , $i \leq n$, is a (possibly empty) conjunction $\delta_{i_1}, \dots, \delta_{i_h}$ of literals (either positive or negative), where for each δ_{i_j} , $i_j \leq i_h$, $\delta_{i_j} \neq \lambda_i$ and $\delta_{i_j} \neq \text{not } \lambda_i$. The Δ_i 's are called the *AND handles* of the cycle. We say that Δ_i is an AND handle for atom λ_i , or, equivalently, an AND handle referring to λ_i .

We say that C has size n and it is even (respectively odd) if $n = 2k$, $k \geq 1$ (respectively, $n = 2k + 1$, $k \geq 0$). For $n = 1$ we have the (odd) self-loop $\lambda_1 \leftarrow \text{not } \lambda_1, \Delta_1$. In what follows, λ_{i+1} will denote $\lambda_{(i+1) \bmod n}$, i.e., $\lambda_{n+1} = \lambda_1$.

A *positive cycle* is similar to a negative cycle, except that we have positive literals λ_i 's in the body of rules instead of negative ones. In the rest of the paper we will consider negative cycles unless differently specified explicitly.

For any cycle C , we will denote by $\text{Composing_atoms}(C)$ the set $\{\lambda_1, \dots, \lambda_n\}$, i.e., the set of atoms *involved* in cycle C . We say that the rules listed above are *involved* in the cycle, or *form* the cycle. In the rest of the paper, sometimes it will be useful to see $\text{Composing_atoms}(C)$ as divided into two subsets, that we indicate as two *kinds* of atoms: the set of the $\text{Even_atoms}(C)$ composed of the λ_i 's with i even, and the set $\text{Odd_atoms}(C)$, composed of the λ_i 's with i odd.

Conventionally, in the rest of the paper C and C_i denote cycles in general, OC and OC_i denote odd cycles, and EC or EC_i denote even cycles.

In the following program for instance, there is an odd cycle that we may call OC_1 , with composing atoms $\{e, f, g\}$ and an even cycle that we may call EC_1 , with composing atoms $\{a, b\}$.

```

—  $OC_1$ 
 $e \leftarrow \text{not } f, \text{not } a$ 
 $f \leftarrow \text{not } g, b$ 
 $g \leftarrow \text{not } e$ 
—  $EC_1$ 
 $a \leftarrow \text{not } b$ 
 $b \leftarrow \text{not } a$ 

```

OC_1 has an AND handle $\text{not } a$ referring to e , and an AND handle b referring to f .

Notice that the sets of atoms composing different cycles are not required to be disjoint. In fact, the same atom may be involved in more than one cycle. The set of atoms composing a cycle can even be a proper subset of the set of atoms composing another cycle, like in the following program, where there is an even cycle EC_1 with composing atoms $\{a, b\}$, since a depends on $\text{not } b$ and b depends on $\text{not } a$, but also an odd cycle OC_1 with composing atom $\{a\}$, since a depends on $\text{not } a$.

```

—  $EC_1$ 
—  $OC_1$ 
 $a \leftarrow \text{not } a, \text{not } b$ 
 $b \leftarrow \text{not } a$ 

```

Here, OC_1 has an AND handle $\text{not } b$ referring to a , but, symmetrically, EC_1 has an AND handle $\text{not } a$ referring to b .

Thus, it may be the case that a handle of a cycle C contains an atom α which is involved in C itself, because α is also involved in some other cycle C_1 .

Definition 5

A rule is called an *auxiliary rule of cycle C* (or, equivalently, *to cycle C*) if it is of this form:

$\lambda_i \leftarrow \Delta$

where $\lambda_i \in \text{Composing_Atoms}(C)$, and Δ is a non-empty conjunction $\delta_{i_1}, \dots, \delta_{i_h}$ of literals (either positive or negative), where for each δ_{i_j} , $i_j \leq i_h$, $\delta_{i_j} \neq \lambda_i$ and $\delta_{i_j} \neq \text{not } \lambda_i$. Δ is called an *OR handle* of cycle C (more specifically, an OR handle for λ_i or, equivalently, and OR handle referring to λ_i).

A cycle may possibly have several auxiliary rules, corresponding to different OR handles. In the rest of this paper, we will call $\text{Aux}(C)$ the set of the auxiliary rules of a cycle C .

In the following program for instance, there is an odd cycle OC_1 with composing atoms $\{c, d, e\}$ and an even cycle EC_1 with composing atoms $\{a, b\}$. The odd cycle has three auxiliary rules.

```

—  $OC_1$ 
 $c \leftarrow \text{not } d$ 
 $d \leftarrow \text{not } e$ 
 $e \leftarrow \text{not } c$ 
—  $\text{Aux}(OC_1)$ 
 $c \leftarrow \text{not } a$ 
 $d \leftarrow \text{not } a$ 
 $d \leftarrow \text{not } b$ 
—  $EC_1$ 
 $a \leftarrow \text{not } b$ 
 $b \leftarrow \text{not } a$ 

```

In particular, we have $\text{Aux}(OC_1) = \{c \leftarrow \text{not } a, d \leftarrow \text{not } a, d \leftarrow \text{not } b\}$.

In summary, a cycle may have some AND handles, occurring in one or more of the rules that form the cycle itself, and also some OR handles, occurring in its auxiliary rules. Cycles and handles can be unambiguously identified on the Extended Dependency Graph (EDG) of the program (Brignoli et al. 1999).

A cycle may also have no AND handles and no OR handles, i.e., no handles at all, in which case it is called *unconstrained*. The following program is composed of unconstrained cycles (in particular, there is an even cycle involving atoms a and b , and an odd cycle involving atom p).

```

—  $EC_1$ 
 $a \leftarrow \text{not } b$ 
 $b \leftarrow \text{not } a$ 
—  $OC_1$ 
 $p \leftarrow \text{not } p$ 

```

Notice that the basic definition of a cycle corresponds to that of a *negative cycle* in the *atom Dependency Graph* as defined in (Fages 1994). However, as discussed in (Costantini 2001), on the dependency graph it is impossible to identify the handles, and there are different programs with different answer set, but the same dependency graph. The handles can be identified unambiguously on the *Extended Dependency Graph* as defined and discussed in (Brignoli et al. 1999) and (Costantini et al. 2002).

4 Canonical programs

In order to analyze the relationship between cycles, handles and consistency, below we introduce a *canonical form* for logic programs. This canonical form is new, and is introduced with the general objective of simplifying the study of formal properties of logic programs under the Answer Set semantics. Rules in *canonical programs* are in a standard format, so as to make definitions and proofs cleaner and easier to read. There is however no loss of generality, since, as proved in the companion paper (Costantini and Proveti 2004), any logic program can be *reduced* to a canonical program, and that stable models of the original program can be easily obtained from the stable models of its canonical counterpart.

Definition 6

A logic program Π is in canonical form (or, equivalently, Π is a canonical program) if it is *WF*-irreducible, and fulfills the following syntactic conditions.

1. Π does not contain positive cycles;
2. every atom in Π occurs both in the head of some rule and in the body of some other (possibly the same) rule;
3. every atom in Π is involved in some cycle;
4. each rule of Π is either involved in a cycle, or is an auxiliary rule of some cycle;
5. each handle of a cycle C consists of exactly one literal, either α or *not* α , where atom α does not occur in C .

Since the above definition requires handles to consist of just one literal, it implies that in a canonical program Π : (i) the body of each rule which is involved in a cycle consists of either one or two literals; (ii) the body of each rule which is an auxiliary rule of some cycle consists of exactly one literal.

Nothing prevents a rule to be at the same time involved in a cycle, and auxiliary to some other cycle. In this case however, the definition requires the rule to have exactly one literal in the body, i.e., the rule cannot have an AND handle.

All definitions and results introduced in the rest of the paper might be rephrased for the general case, but the choice of referring to canonical programs is a significant conceptual simplification that leads without loss of generality to a more readable and intuitive formalization. Notice for instance that in canonical programs the problem of consistency arises only in cycles containing an odd number of rules, since cycles do not have non-negated composing atoms.

Although for a detailed discussion we refer to (Costantini and Proveti 2004), it is important to recall that canonical programs are *WF*-irreducible. For instance, the program

$$\begin{aligned} p &\leftarrow \text{not } p, q \\ q &\leftarrow \text{not } q, p \end{aligned}$$

may look canonical, while it is not, since it has a non-empty well-founded model $\langle \emptyset; \{p, q\} \rangle$. In particular, since there are no undefined atoms, the set of true atoms of the well-founded model (in this case $\emptyset$) coincides (as it is well-known) with the unique stable model.

Similarly, the program

$$\begin{aligned} q &\leftarrow \text{not } q \\ q &\leftarrow p \end{aligned}$$

may look canonical, while it is not, since it has a non-empty well-founded model $\langle \emptyset; \{p\} \rangle$. Atom q is undefined. The corresponding canonical program is $q \leftarrow \text{not } q$ that, like the original program, has no stable models. The second rule is dropped by canonization, since its condition is false w.r.t. the well-founded model. The program

$$\begin{aligned} q &\leftarrow p \\ p &\leftarrow \text{not } r \\ r &\leftarrow \text{not } q \end{aligned}$$

is not canonical because atom p is not involved in any cycle. In fact, in order to be involved in a cycle an atom must occur in the head of some rule but, also, its negation must occur in the body of some, possibly different, rule. Here, atom p forms an (inessential) intermediate step between the two atoms q and r that actually form a cycle. The canonical form of the program is $q \leftarrow \text{not } r, r \leftarrow \text{not } q$. Given the stable models $\{q\}$ and $\{r\}$ of the canonical program, the stable models $\{p, q\}$ and $\{r\}$ of the original program can be easily obtained, since the truth value of p directly depends on that of r .

In the following, let the program at hand be a logic program Π in canonical form, unless differently specified explicitly. Let $C_1, \dots, C_w$ be all the cycles occurring in Π (called *the composing cycles* of Π). Whenever we will refer to C, C_1, C_2, C_i etc. we implicitly assume that these are cycles occurring in Π .

5 Active handles

In this section we make some preliminary steps toward establishing a connection between syntax (cycles and handles) and semantics (consistency of the program). Truth or falsity of the atoms occurring in the handles of a cycle (w.r.t. a given set of atoms) affects truth/falsity of the atoms involved in the cycle. As we discuss at length in the rest of the paper, this creates the conditions for stable models to exist or not, and these conditions can be checked on the Cycle Graph of the program. This idea is formalized in the following definitions of *active handles*, that will be frequently used in the rest of the paper.

Definition 7

Let $\mathcal{I}$ be a set of atoms. An AND handle Δ of cycle C is *active* w.r.t. $\mathcal{I}$ if it is false w.r.t. $\mathcal{I}$. We say that the rule where the handle occurs has an *active AND handle*. An OR handle Δ of cycle C is *active* w.r.t. $\mathcal{I}$ if it is true w.r.t. $\mathcal{I}$. We say that the rule where the handle occurs has an *active OR handle*.

Assume that $\mathcal{I}$ is a model. We can make the following observations. (i) The head λ of a rule ρ with an active AND handle is not required to be true in $\mathcal{I}$. (ii) The head of a rule $\lambda \leftarrow \Delta$ where Δ is an active OR handle is necessarily true in $\mathcal{I}$: since the body is true, the head λ must also be true.

Observing which are the active handles of a cycle C gives relevant indications about whether a set of atoms $\mathcal{I}$ is a stable model.

Consider for instance the following program:

```

—  $OC_1$ 
 $p \leftarrow not\ p, not\ a$ 
—  $EC_1$ 
 $a \leftarrow not\ b$ 
 $b \leftarrow not\ a$ 
—  $OC_2$ 
 $q \leftarrow not\ q$ 
—  $Aux(OC_2)$ 
 $q \leftarrow f$ 
—  $EC_2$ 
 $e \leftarrow not\ f$ 
 $f \leftarrow not\ e$ 

```

The sets of atoms $\{a, f, q\}$, $\{a, e, q\}$, $\{b, p, f, q\}$ $\{b, p, e, q\}$ are minimal models. Consider the set of atoms $\{a, f, q\}$: both the AND handle $not\ a$ of cycle $p \leftarrow not\ p, not\ a$ and the OR handle f of cycle $q \leftarrow not\ q$ are active w.r.t. this set of atoms. $\{a, f, q\}$ is a stable model, since atom p is forced to be false, and atom q is forced to be true, thus avoiding the inconsistencies. In all the other minimal models instead, one of the handles is not active. I.e., either literal $not\ a$ is true, and thus irrelevant in the context of a rule which is inconsistent, or literal f is false, thus leaving the inconsistency on q . These minimal models are in fact not stable.

In conclusion, the example suggests that for a minimal model $\mathcal{M}$ to be stable, each odd cycle must have an active handle. Formally:

Theorem 1

Let Π be a program, and let $\mathcal{M}$ be a minimal model of Π . $\mathcal{M}$ is a stable model only if each odd cycle OC_i occurring in Π has an active handle w.r.t. $\mathcal{M}$.

Proof

Since $\mathcal{M}$ is stable, for each $A \in \mathcal{M}$ there exists a rule in Π with head A and body which is true w.r.t. $\mathcal{M}$, i.e., a rule which *supports* A . Let $x \bmod y$ be (as usual) the remainder of the integer division of x by y .

Assume that $\mathcal{M}$ is stable, but there is an odd cycle without active handles, composed of atoms $\lambda_1, \dots, \lambda_n$, where n is odd. Take a λ_i , and assume first that $\lambda_i \in \mathcal{M}$. Since there is no active OR handle, each λ_i can possibly be supported only by the corresponding rule in the cycle. By definition of cycle, this rule has the form:

$$\lambda_i \leftarrow not\ \lambda_{(i+1) \bmod n}, \Delta_i$$

Since there are no active AND handles, then all Δ 's are true w.r.t. $\mathcal{M}$. For λ_i to be sup-

ported, *not* $\lambda_{(i+1) \bmod n}$ should be true as well, i.e., $\lambda_{(i+1) \bmod n}$ should be false. The rule for $\lambda_{(i+1) \bmod n}$ has the form:

$$\lambda_{(i+1) \bmod n} \leftarrow \text{not } \lambda_{(i+2) \bmod n}, \Delta_{(i+1) \bmod n}$$

Since $\Delta_{(i+1) \bmod n}$ is true w.r.t. $\mathcal{M}$, for $\lambda_{(i+1) \bmod n}$ to be false, *not* $\lambda_{(i+2) \bmod n}$ should be false as well, i.e., $\lambda_{(i+2) \bmod n}$ should be true. By iterating this reasoning, $\lambda_{(i+3) \bmod n}$ should be false, etc. In general, $\lambda_{(i+k) \bmod n}$ should be false w.r.t. $\mathcal{M}$ with k odd, and true with k even. Then, since the number n of the composing atoms is odd, $\lambda_{(i+n) \bmod n}$ should be false w.r.t. $\mathcal{M}$, but $\lambda_{(i+n) \bmod n} = \lambda_i$, which is a contradiction. Assume now that $\lambda_i \notin \mathcal{M}$. Then, *not* λ_i is true w.r.t. $\mathcal{M}$, and thus, since the corresponding AND handle is not active, $\lambda_{(i-1) \bmod n}$ is supported and should belong to $\mathcal{M}$. Consequently, we should have $\lambda_{(i-2) \bmod n} \notin \mathcal{M}$. In general, $\lambda_{(i-k) \bmod n}$ should be true w.r.t. $\mathcal{M}$ with k odd, and false with k even. Then, since the number n of the composing atoms is odd, $\lambda_{(i-n) \bmod n}$ should be true w.r.t. $\mathcal{M}$, but $\lambda_{(i-n) \bmod n} = \lambda_i$, which is again a contradiction. $\square$

Another thing that the above example shows is that the stable model $\{a, f, q\}$ of the overall program is actually the union of the stable model $\{a\}$ of the program fragment $OC_1 \cup EC_1$ and of the stable model $\{f, q\}$ of the program fragment $OC_2 \cup Aux(OC_2) \cup EC_2$. This is not by chance, and in the next sections we will study how to relate the existence of stable models of the overall program to the existence of stable models of the composing cycles.

In order to do so, some preliminary definitions about handles are in order. It is useful to collect the set of handles of a cycle into a set, where however each handle is annotated so as to keep track of its *kind*. I.e., we want to remember whether a handle is an OR handle or an AND handle of the cycle.

Definition 8

Given cycle C , the set H_C of the handles of C is defined as follows, where $\beta \in Composing_Atoms(C)$:

$$H_C = \{(\Delta : AND : \beta) \mid \Delta \text{ is an AND handle of } C \text{ referring to } \beta\} \cup \{(\Delta : OR : \beta) \mid \Delta \text{ is an OR handle of } C \text{ referring to } \beta\}$$

Whenever we need not care about β we shorten $(\Delta : K : \beta)$ as $(\Delta : K)$, $K = \text{AND/OR}$. We call “handles” the expressions in both forms, and whenever necessary we implicitly shift from one form to the other one. Informally, we will say for instance “the OR (resp. AND) handle Δ of β ” meaning $(\Delta : OR : \beta)$ (resp. $(\Delta : AND : \beta)$).

In general however the indication of β is necessary. In fact, different atoms of a cycle may have handles with the same Δ , but although active/not active at the same time, they may affect the existence of stable models differently. Take for instance the following program with the indication of the composing cycles:

$\text{--- } OC_1$
 $q \leftarrow \text{not } q, e$
 $q \leftarrow \text{not } f$
 $\text{--- } OC_2$
 $a \leftarrow \text{not } b, \text{not } e$
 $b \leftarrow \text{not } c, \text{not } f$
 $c \leftarrow \text{not } a, \text{not } e$
 $\text{--- } OC_3$
 $p \leftarrow \text{not } p, \text{not } e$
 $\text{--- } EC_1$
 $e \leftarrow \text{not } f$
 $f \leftarrow \text{not } g$

we have $H_{OC_1} = \{(e : \text{AND} : q), (\text{not } f : \text{OR} : q)\}$, $H_{OC_2} = \{(\text{not } e : \text{AND} : a), (\text{not } f : \text{AND} : b), (\text{not } e : \text{AND} : c)\}$, $H_{OC_3} = \{(\text{not } e : \text{AND} : p)\}$, $H_{EC_1} = \emptyset$. Handle $(\text{not } e : \text{AND})$ occurs several times, even twice in cycle OC_2 , referring to different atoms. Notice that the same literal Δ may occur both in AND and OR handles. E.g., $\text{not } f$ occurs both in an AND handle (of OC_2) and in an OR handle (of OC_1). Notice also that the same atom α may appear in literals α and $\text{not } \alpha$ that occur in different handles. E.g., f occurs in an OR handle of OC_1 , and $\text{not } f$ occurs in an AND handle of OC_2 .

Given any subset Z of H_C , it is useful to identify the set of atoms occurring in the handles belonging to Z .

Definition 9

Let $Z \subseteq H_C$. The set of the atoms occurring in the handles belonging to Z is defined as follows.

$$\begin{aligned} \text{Atoms}(Z) = & \{ \alpha \mid (\alpha : K) \in Z \} \cup \\ & \{ \alpha \mid (\text{not } \alpha : K) \in Z \} \end{aligned}$$

If for instance we take $Z = H_{OC_1}$, we have $\text{Atoms}(H_{OC_1}) = \{e, f\}$.

Given any subset Z of H_C , it is useful to state which are the atoms that are required to be true, in order to make all the handles in Z active (implicitly, to this aim all the other atoms are required to be false).

Definition 10

Let $Z \subseteq H_C$. The set of atoms $\text{ActivationAt}_C(Z) \subseteq \text{Atoms}(Z)$ is defined as follows.

$$\begin{aligned} \text{ActivationAt}_C(Z) = & \{ \alpha \mid (\alpha : \text{OR}) \in Z \} \cup \\ & \{ \alpha \mid (\text{not } \alpha : \text{AND}) \in Z \} \end{aligned}$$

If for instance we take $Z = H_{OC_3}$, we have $\text{ActivationAt}(H_{OC_3}) = \{e\}$.

Vice versa, any subset V of $\text{Atoms}(H_C)$ corresponds to a subset of the handles of C that become active, if atoms in V are true.

Definition 11

Let $V \subseteq \text{Atoms}(H_C)$.

$$\begin{aligned} \text{Active}_C(V) = & \{(\Delta : \text{AND}) \mid \Delta = \text{not } \alpha, \alpha \in V\} \cup \\ & \{(\Delta : \text{OR}) \mid \Delta = \alpha, \alpha \in V\} \end{aligned}$$

If for instance we take $V = \{e\}$ for cycle OC_3 , we have $\text{Active}_{OC_3}(\{e\}) = \{(\text{not } e : \text{AND})\}$.

Finally, it is useful to introduce a short notation for the union of different sets of rules.

Definition 12

Let $I_1, \dots, I_q$ be sets of rules. As a special case, some of the I_j 's can be sets of atoms, where each atom $\beta \in I_j$ is understood as a fact $\beta \leftarrow$. By $I_1 + \dots + I_q$ we mean the program consisting of the union of all the rules belonging to $I_1, \dots, I_q$.

6 Cycle, handles and existence of stable models

In this and the following sections we proceed further toward a framework that relates cycles, handles and active handles to the existence of stable models. This relation is far from obvious, as demonstrated by the following simple program.

— OC_1
 $p \leftarrow \text{not } p, \text{not } a$
 — EC_1
 $a \leftarrow \text{not } b$
 $b \leftarrow \text{not } a$
 — OC_2
 $q \leftarrow \text{not } q, \text{not } b$

In this case, we have only one even cycle, and we might consider the program fragments: (i) $OC_1 \cup EC_1$ with stable model $\{a\}$, based on the active handle $\text{not } a$; (ii) $OC_2 \cup EC_1$ with stable model $\{b\}$, based on the active handle $\text{not } b$. Unfortunately, the union $\{a, b\}$ of the stable models of the two program fragments is a minimal model but it is not stable. In fact, neither atom a nor atom b is supported. This is because the unconstrained even cycle EC_1 , taken per se, has stable models $\{a\}$ and $\{b\}$, which are alternative and cannot be merged: the rules of this cycle in fact state that a holds if b does not hold, and vice versa. Thus, EC_1 cannot provide active handles for both the odd cycles.

Then, if we want to check whether a minimal model is stable, we not only have to check that every odd cycle has an active handle w.r.t. that model, but also that these handles do not enforce contradictory requirements on the even cycles. We can try to build a stable model of the overall program out of the stable models of the composing cycles, taking however care of avoiding inconsistencies on the handles.

Consider a cycle C_i occurring in Π together with its auxiliary rules, i.e., consider the set of rules $C_i + \text{Aux}(C_i)$ and take it as an independent program. Notice that this program is not canonical, since there are atoms that do not appear in the conclusion of rules: these are exactly the atoms occurring in the handles of C_i , i.e., the atoms in $\text{Atoms}(H_{C_i})$. Take a set $X_i \subseteq \text{Atoms}(H_{C_i})$, and assume to add atoms in X_i as facts to $C_i + \text{Aux}(C_i)$.

Definition 13

Let C_i be a cycle. Let $X_i \subseteq H_{C_i}$. The general logic program $C + Aux(C) + X_i$ is called an *extended cycle* of C_i corresponding to X_i .

Depending on the active handles $Active_{C_i}(X_i)$ corresponding to X_i , the extended cycle $C_i + Aux(C_i) + X_i$ may or may not be consistent.

Definition 14

Let C_i be a cycle occurring in Π . We say that C_i has *partial stable models* if $\exists X_i \subseteq Atoms(H_{C_i})$ such that the corresponding extended cycle $C_i + Aux(C_i) + X_i$ is consistent. Given a stable model S_{C_i} of $C_i + Aux(C_i) + X_i$, the set X_i is called a *positive base* for S_{C_i} , while the set $X_i^- = Atoms(H_{C_i}) \setminus X_i$ is called a *negative base* for S_{C_i} . The couple of sets $\langle X_i, X_i^- \rangle$ is called a *base* for S_{C_i} . We say that the S_{C_i} 's are *partial stable models* for Π relative to C_i .

Atoms in X_i are added as facts in order to simulate that we deduce them true in some other part of the program. Symmetrically, atoms in X_i^- are supposed not to be concluded true anywhere in the program. The positive base X_i may be empty: in this case, all the atoms occurring in the handles are supposed to be false. Clearly, there may be no partial stable models relative to a cycle C_i or there may be several ones. However, partial stable models of cycles are related to stable models of the overall program.

Lemma 1

Let Π be a program, C_i be one of its composing cycle, and $\mathcal{I}$ be a stable model of Π . Let $X_i = \mathcal{I} \cap Atoms(H_{C_i})$. Then, the restriction S_i of $\mathcal{I}$ to the atoms involved in the extended cycle $P_i = C_i + Aux(C_i) + X_i$ is a partial stable model of P_i .

Proof

Notice that all non-unit rules of P_i are also rules of Π . Notice also that for every atom α occurring in P_i as the head of a non-unit rule, P_i contains all rules of Π with head α : as Π is canonical, these rules are by Definition 6 either in C_i or in $Aux(C_i)$. Assume that S_i is not a stable model of P_i . This means that, after applying the reductions specified in Definition 2, we obtain a positive program $P_i^{S_i}$ where either (i) there exists atom $\alpha \in S_i$ that is not a consequence of $P_i^{S_i}$ or (ii) there exists atom β which is a consequence of $P_i^{S_i}$, but $\beta \notin S_i$. In situation (i), it means that we have canceled all rules with head α , because they contain a negative literal which is false w.r.t. S_i . But, as we have included in S_i all the atoms of $\mathcal{I}$ that occur either in C_i or in its OR handles, this rules would have been canceled w.r.t. Π as well, and thus $\mathcal{I}$ could not be stable. In situation (ii), there is some rule that we would have canceled w.r.t. P and has not been canceled for P_i , i.e, there is a literal *not* α which is false w.r.t. $\mathcal{I}$ and true w.r.t. S_i . But, if $\alpha \in \mathcal{I}$ and α occurs in P_i , then $\alpha \in S_i$ by hypothesis, and thus (ii) cannot be the case as well. $\square$

Once we get partial stable models of the composing cycles, we can try to put them together in order to obtain stable models of the whole program. Of course, we will try to obtain a stable model of the overall program by taking one partial stable model for each cycle, and considering their union. This however will work only if the partial stable models assign truth values to atoms in a compatible way.

Definition 15

Consider a collection $\mathcal{S} = S_1, \dots, S_w$ of partial stable models for Π , relative to its composing cycles $C_1, \dots, C_w$, each S_i with base $\langle X_i, X_i^- \rangle$. We say that $S_1, \dots, S_w$ are *compatible* or, equivalently, that $\mathcal{S}$ is a *compatible set of partial stable models* whenever the following conditions hold:

1. $\forall j, k \leq w, X_j \cap X_k^- = \emptyset$;
2. $\forall j \leq w, \forall A \in X_j, \exists h \neq j$ such that $A \in S_h$, and $A \notin X_h$;
3. $\forall j \leq w, \forall B \in X_j^-, \nexists t \leq w$ such that $B \in S_t$.

Condition (1) states that the bases of compatible partial stable models cannot assign opposite truth values to any atom. Condition (2) ensures that, if an atom A is supposed to be true in the base of some cycle C_j , it must be actually concluded true in some other cycle C_h . Notice that “concluded” does not mean “assumed”, and thus A must occur in the partial stable model S_h of C_h , without being in its set of assumptions X_h . Condition (3) ensures that, if an atom is supposed to be false in the base of some cycle, it cannot be concluded true in any of the other cycles.

The following result formally states the connection between the stable models of Π , and the partial stable models of its cycles.

Theorem 2

Let Π be a program with composing cycles $C_1, \dots, C_w$ and $\mathcal{I}$ be a set of atoms. $\mathcal{I}$ is a stable model of Π if and only if $\mathcal{I} = \bigcup_{i \leq w} S_i$ where each S_i is a partial stable model for C_i and $\mathcal{S} = S_1, \dots, S_w$ is a compatible set of partial stable models.

Proof

Suppose that $\mathcal{I}$ is a stable model for Π . Let $C_i, i \leq w$, be any of the composing cycles of Π . Let $X_i = \mathcal{I} \cap \text{Atoms}(H_{C_i})$, which means that X_i is the set of the atoms of the handles of C_i which are true w.r.t. $\mathcal{I}$, and $X_i^- = \text{Atoms}(H_{C_i}) \setminus X_i$. Let S_i be the restriction of $\mathcal{I}$ to the atoms involved in the extended cycle $P_i = C_i + \text{Aux}(C_i) + X_i$. By Lemma 1 S_i is a stable model for P_i . Then, it remains to prove that $S_1, \dots, S_w$ form a compatible set of partial stable models. Condition 1 of Definition 15 holds because by construction we put in X_i all the atoms occurring in the handles of C_i that are true w.r.t. $\mathcal{I}$: should they occur in the handles of some other cycle C_j they should be in X_j , and not in X_j^- . For condition 2, notice that atoms in X_i do not occur in the head of the rules of C_i . Since they occur in $\mathcal{I}$, they must have been derived by means of the rules of some other cycle C_j , and thus they occur in S_j . For condition 3, it is sufficient to notice that we put in the X_j^- ’s atoms that are not in $\mathcal{I}$, and consequently are not in the S_i ’s.

Vice versa, let us consider a compatible set $\mathcal{S} = S_1, \dots, S_w$ of partial stable models for the cycles in Π . Notice that Π itself corresponds to the union of the cycles and of their auxiliary rules, i.e., $\Pi = \bigcup_{i \leq w} C_i + \text{Aux}(C_i)$.

Let us first show that $\mathcal{I} = \bigcup_{i \leq w} S_i$ is a stable model of the program $\Pi_L = \bigcup_{i \leq w} C_i + \text{Aux}(C_i) + X_i$, which is a superset of Π . In fact, each S_i satisfies the stability condition on the rules of the corresponding extended cycle, and, since they form a compatible set, by

conditions (1) and (3) of Definition 15 no atom which is in the negative base of any of the S_i 's, is concluded true in some other S_j . Therefore, $\mathcal{I}$ is a stable model of Π_L .

In order to obtain Π from Π_L , we have to remove the positive bases of cycles, which are the unit rules corresponding to the X_i 's. By condition (2) however, in a set of compatible partial stable models every atom $A \in X_i$ is concluded true in some S_j , $i \neq j$, i.e., in the partial stable model of some other cycle. This implies that $\mathcal{I}$ satisfies the stability condition also after X_i 's have been removed: then, $\mathcal{I}$ is a stable model for Π . $\square$

Each stable model S of Π corresponds to a different choice of the X_i 's, i.e., of the active handles of cycles.

The above result is of theoretical interest, since it sheds light on the connection between stable models of a program and stable models of its sub-parts. It may also contribute to any approach to modularity in software development under the stable model semantics.

From Theorem 1 and Theorem 2 we can argue that for checking whether a logic program has stable models (and, possibly, for finding these models) one can do the following.

- (i) Divide program Π into pieces, of the form $C_i + Aux(C_i)$, and check whether every odd cycle has handles; if not, then the program is inconsistent.
- (ii) For every cycle C_i with handles, find the sets X_i that make the subprogram $C_i + Aux(C_i)$ consistent, and find the stable models S_{C_i} of each $C_i + Aux(C_i) + X_i$. Notice that in the case of unconstrained even cycles, H_{C_i} is empty, and we have two stable models, namely $M_{C_i}^1 = \text{Even_atoms}(C_i)$ and $M_{C_i}^2 = \text{Odd_atoms}(C_i)$.
- (iii) Check whether there exists a collection of X_i 's, one for each cycle, such that the corresponding S_{C_i} 's form a compatible collection of partial stable models for Π : in this case the program is consistent, and its stable model(s) can be obtained as the union of the S_{C_i} 's.

To show how the method works, consider for instance the following program.

```

—  $OC_1$ 
 $q \leftarrow not\ q$ 
—  $Aux(OC_1)$ 
 $q \leftarrow f$ 
—  $OC_2$ 
 $p \leftarrow not\ p, not\ f$ 
—  $EC_1$ 
 $e \leftarrow not\ f$ 
 $f \leftarrow not\ e$ 

```

It can be seen as divided into the following parts, each one corresponding to $C_i + Aux(C_i)$ for cycle C_i . The first part is composed of odd cycle OC_1 , with an auxiliary rule (OR handle):

```

 $q \leftarrow not\ q$ 
 $q \leftarrow f$ 

```

The second part is composed of odd cycle OC_2 , without auxiliary rules but with an AND handle:

$p \leftarrow \text{not } p, \text{not } f$

The third part is composed of the unconstrained even cycle EC_1 :

$e \leftarrow \text{not } f$

$f \leftarrow \text{not } e$

OC_1 in itself is inconsistent, but if we take $X_{OC_1} = \{f\}$ (and $X_{OC_1}^- = \emptyset$) we get the partial stable model $\{f, q\}$: the active OR handle forces q to be true. Similarly, if we take for OC_2 $X_{OC_2} = \{f\}$, we get the partial stable model $\{f\}$: the active AND handle forces p to be false. Cycle EC_1 is consistent, with partial stable models $\{e\}$ and $\{f\}$. If we now select the partial stable model $\{f\}$, we get a compatible set of partial stable models thus obtaining the stable model $\{f, q\}$ for the overall program. Instead, the partial stable model $\{e\}$ for EC_1 does not serve to the purpose of obtaining a stable model for the overall program, since condition 2 of Definition 15 is not fulfilled. This in particular means that atom f , which is in the positive base of both the odd cycles, is not concluded true in this partial stable model. Therefore, the handles of the odd cycles are not active and no overall consistency can be achieved.

Take now this very similar program, that can be divided into cycles analogously.

— OC_1

$q \leftarrow \text{not } q$

— $Aux(OC_1)$

$q \leftarrow f$

— OC_2

$p \leftarrow \text{not } p, \text{not } e$

— EC_1

$e \leftarrow \text{not } f$

$f \leftarrow \text{not } e$

The difference is that OC_2 has AND handle $\text{not } e$ (instead of $\text{not } f$). With base $X_{OC_2} = \{e\}$ we get partial stable model $\{e\}$. To fulfill condition 2 of Definition 15, we should select partial stable model $\{e\}$ of EC_1 . Unfortunately however, since OC_1 is consistent only if we take $X_{OC_1} = \{f\}$, we should at the same time choose the other partial stable model $\{f\}$ of EC_1 . Thus, no choice can be made for EC_1 so as to make this program consistent.

With the aim of developing effective software engineering tools and more efficient algorithms for computing stable models, syntactic conditions for the existence of stable models are in order. In the ongoing, we will use the above results as the basis for defining necessary and sufficient syntactic conditions for consistency.

7 Handle assignments and admissibility

In previous sections we have discussed how to split a stable model of a program into a compatible set of partial stable models of the cycle. However, we have not formalized a method for selecting bases for the cycles so as to ensure that all cycles have partial stable models, and that they form a compatible collection. To this aim, in this section we define syntactic condition that specify how active handles affect consistency of extended cycles.

Assume that atom α occurs in some handle Δ of a cycle. Then, it may possibly occur in the positive or negative base of that cycle for forming a partial stable model that may be a part of a compatible collection. In this case: if α occurs in the positive base, then by condition 2 of Definition 15 it must be concluded true in some other cycle; if instead α occurs in the negative base, then by condition 3 of Definition 15 it must not be concluded true in any other cycle. Notice that the cycles where it is possible to derive an atom α are the cycles α is involved in, which are the cycles the handle Δ comes from, or equivalently the *source cycles* of the handle.

Definition 16

A handle ($\Delta : K$) of cycle C_1 , $\Delta = \alpha$ or $\Delta = \text{not } \alpha$ comes from source cycle C_2 if $\alpha \in \text{Composing_atoms}(C_2)$.

Handles in H_C are called the *incoming handles* of C . The same handle of a cycle C may come from different cycles, and may refer to different atoms of C . For instance, in the program below we have:

```

— OC1
a ← not b, not f
b ← not c
c ← not a, not f
b ← g
— EC1
f ← not g
g ← not f
— EC2
f ← not h
h ← not f

```

handle ($\text{not } f : \text{AND}$) of OC_1 comes from both EC_1 and EC_2 , and refers to two different atoms in OC_1 , namely a and c ; handle ($g : \text{OR}$) of OC_1 comes from EC_1 , and refers to atoms b .

The following definition completes the terminology by identifying the atoms occurring in handles coming from C .

Definition 17

Given a cycle C , the set of the atoms involved in C that occur in the handles of some other cycle is defined as follows:

$$\text{Out_handles}(C) = \{\beta \mid \beta \in \text{Composing_Atoms}(C) \wedge \exists C_1 \neq C \text{ such that } \beta \in \text{Atoms}(H_{C_1})\}$$

In the above program for instance, we have $\text{Out_handles}(EC_1) = \{f, g\}$ and $\text{Out_handles}(EC_2) = \{f\}$. Notice that, according to the definition, $h \notin \text{Out_handles}(EC_2)$, because h does not occur in any other cycle.

For an handle to be active w.r.t. a set of atoms, we must have the following. (i) If the corresponding atom α is required to be true, then it must be concluded true (by means of a

supporting rule) in *at least one* of the cycles the handle comes from, which implies α to be concluded true in *all* the *extended* cycles it is involved into: in fact, the rule that makes α true is an auxiliary rule for all these cycles. (ii) If the corresponding atom α is required to be false, then it must be concluded false in *all* the (extended) cycles it comes from.

This is illustrated by the following example:

— OC_1
 $p \leftarrow \text{not } p, \text{not } c$
 — OC_2
 $c \leftarrow \text{not } d$
 $d \leftarrow \text{not } e$
 $e \leftarrow \text{not } c, f$
 — EC_1
 $f \leftarrow \text{not } g$
 $g \leftarrow \text{not } f$
 — EC_2
 $f \leftarrow \text{not } h$
 $h \leftarrow \text{not } f$

The extended cycles are:

— OC_1
 $p \leftarrow \text{not } p, \text{not } c$

with no auxiliary rules, $\text{Out_handles}(EC_1) = \emptyset$, $H_{OC_1} = \{\text{not } c : \text{AND} : p\}$, $\text{Atoms}(H_{OC_1}) = \{c\}$ and unique partial stable model $\{c\}$ obtained by choosing $X_{OC_1} = \{c\}$;

— OC_2
 $c \leftarrow \text{not } d$
 $d \leftarrow \text{not } e$
 $e \leftarrow \text{not } c, f$

with no auxiliary rules, $\text{Out_handles}(OC_2) = \{c\}$, $H_{OC_2} = \{f : \text{AND} : e\}$, $\text{Atoms}(H_{OC_2}) = \{f\}$ and unique partial stable model $\{d\}$ obtained by choosing positive base $X_{OC_2} = \emptyset$, $X_{OC_1}^- = \{f\}$;

— $EC_1 + \text{Aux}(EC_1)$
 $f \leftarrow \text{not } g$
 $g \leftarrow \text{not } f$
 $f \leftarrow \text{not } h$

with $\text{Out_handles}(EC_1) = \{f\}$, $H_{EC_1} = \{\text{not } h : \text{OR} : f\}$, $\text{Atoms}(H_{EC_1}) = \{h\}$. Notice that if we take $X_{EC_1} = \{h\}$ this makes f not derivable via the auxiliary rule, while however f is still derivable via the corresponding rule involved in the cycle. Then, if we consider the two partial stable models $\{f\}$ and $\{g\}$ of EC_1 , we see that: the former one can be obtained either by choosing either $X_{EC_1} = \emptyset$ or $X_{EC_1} = \{h\}$; the latter one instead requires $X_{EC_1} = \{h\}$, so as to allow f to be false.

— $EC_2 + Aux(EC_2)$

$f \leftarrow not\ h$

$h \leftarrow not\ f$

$f \leftarrow not\ g$

with $Out_handles(EC_2) = \{f\}$, $H_{EC_2} = \{not\ g : OR : f\}$, $Atoms(H_{EC_2}) = \{g\}$ and two partial stable models $\{f\}$ and $\{h\}$. The former one can be obtained either by choosing $X_{EC_2} = \emptyset$ or, also, $X_{EC_1} = \{g\}$. The latter one requires $X_{EC_2} = \{g\}$, so as to allow f to be false.

Unfortunately, the overall program turns out to have no stable model, because: for obtaining the partial stable model of OC_2 , f must be concluded false so to make the unique AND handle active. Both EC_1 and EC_2 actually admit a partial stable model where f is false. Thus, for the fragment $EC_1 + EC_2 + OC_2$ we may construct the unique wider partial stable model $\{g, h, d\}$. However, this fails to make the handle of OC_1 active, and therefore a stable model for the program cannot be obtained.

Assume to replace OC_1 with OC'_1

— OC'_1

$p \leftarrow not\ p, not\ d$

where $H_{OC'_1} = \{not\ d : AND : p\}$, $Atoms(H_{OC'_1}) = \{d\}$ and there is a unique partial stable model $\{d\}$ obtained by choosing positive base $X_{OC_1} = \{d\}$. In this case, $\{g, h, d\}$ would be a stable model for the overall program.

Assume to add the cycles:

— OC_3

$q \leftarrow not\ q, d$

— OC_4

$r \leftarrow not\ r$

$r \leftarrow not\ d.$

The resulting program cannot be consistent. On the one hand in fact, OC'_1 and OC_3 have unique AND handles $not\ d$ and d respectively, that cannot be active at the same time. On the other hand, OC'_1 has an AND handle $not\ d$ while OC_4 has an OR handle $not\ d$, and also in this case these handles cannot be active at the same time.

Below we establish the formal foundations of the kind of reasoning that we have informally proposed up to now. Some more definitions about handles are needed.

Definition 18

The handles $(\Delta : AND)$ and $(\Delta : OR)$ are called *opposite handles*. Given a handle h , we will indicate its opposite handle with h^- .

Definition 19

The handles $(\Delta_1 : K)$ and $(\Delta_2 : K)$ are called *contrary handles* if $\Delta_1 = \alpha$ and $\Delta_2 = not\ \alpha$. Given a handle h , we will indicate its contrary handle with h^n .

Whenever either contrary or opposite pairs of handles occur in a program, even for different β 's, if one is active w.r.t. a given set of atoms then the other one is not active, and vice versa.

Definition 20

The handles $(\Delta_1 : K1)$ and $(\Delta_2 : K2)$ are called *sibling handles* if $K1 \neq K2$, and $\Delta_1 = \alpha$ and $\Delta_2 = \text{not } \alpha$. Given a handle h , we will indicate its sibling handle with h^s .

Whenever sibling pairs of handles occur in a program, even for different β 's, if one is active if one is active w.r.t. a given set of atoms then the other one is active as well.

Taken for instance atom α , we have:

- $(\alpha : AND)$ and $(\alpha : OR)$ are opposite handles;
- $(\text{not } \alpha : AND)$ and $(\text{not } \alpha : OR)$ are opposite handles;
- $(\alpha : AND)$ and $(\text{not } \alpha : AND)$ are contrary handles;
- $(\alpha : OR)$ and $(\text{not } \alpha : OR)$ are contrary handles;
- $(\alpha : OR)$ and $(\text{not } \alpha : AND)$ are sibling handles;
- $(\alpha : AND)$ and $(\text{not } \alpha : OR)$ are sibling handles;

Finally, we introduce the definition of *handle assignment*, which is a consistent hypothesis on (some of) the handles of a cycle C . Namely, it is a quadruple composed of the following sets. IN_C^A contains the incoming handles which are assumed to be active. From IN_C^A one can immediately derive a corresponding assumption on X_C . In particular, $X_C = \text{ActivationAt}_C(IN_C^A)$, i.e. it is exactly the set of the atoms that make the handles in IN_C^A active. Vice versa, IN_C^N contains the incoming handles which are assumed to be not active. Handles of C which are not in $IN_C^A \cup IN_C^N$ can be either active or not active, but their status is either unknown or irrelevant in the context where the handle assignment is used.

OUT_C^+ is the set of out-handles which are required to be concluded true. This in order to make some handle of some other cycle active, as we have seen in the example above. Similarly, OUT_C^- is the set of the out-handles which are required to be concluded false, for the same reason. Of course, the OUT_C 's must be disjoint, since no atom can be required to be simultaneously true and false.

Definition 21

A *basic handle assignment* to (or for) cycle C is a quadruple of sets

$$\langle IN_C^A, IN_C^N, OUT_C^+, OUT_C^- \rangle$$

where the (possibly empty) composing sets are such that:

$$IN_C^A \cup IN_C^N \subseteq H_C;$$

$$IN_C^A \cap IN_C^N = \emptyset;$$

neither IN_C^A and IN_C^N contain pairs of either opposite or contrary handles;

$$OUT_C^+ \cup OUT_C^- \subseteq \text{Out_handles}(C);$$

$$OUT_C^+ \cap OUT_C^- = \emptyset.$$

For short, when talking of both IN_C^A and IN_C^N we will say “the IN_C ’s”. A handle assignment will be called *trivial* (resp. non-trivial) if $OUT_C^+ = OUT_C^- = \emptyset$, i.e., whenever there is no requirement on the out-handles of C .

If IN_C^A is empty, there are two possible situations. (i) $H_C = \emptyset$, i.e., the cycle is unconstrained. (ii) $H_C \neq \emptyset$ but no active incoming handle is assumed: in this case, we say that the cycle is *actually unconstrained* w.r.t. this handle assignment. A handle assignment will be called *effective* (w.r.t. non-effective) whenever $IN_C^A \neq \emptyset$.

We have to cope with the relationship between opposite, contrary, and sibling handles, whenever they should occur in the same cycle C .

Definition 22

Let: h and h^- be a pair of opposite handles; h and h^n be a pair of contrary handles; and h and h^s be a pair of sibling handles. A *complete handle assignment*, or simply a *handle assignment*, to cycle C is a basic handle assignment to C where, for each pair of opposite, contrary or sibling handles the occur in C , the following conditions hold:

$h \in IN_C^A$ if and only if $h^- \in IN_C^N$;

$h \in IN_C^A$ if and only if $h^n \in IN_C^N$;

then either $h, h^s \in IN_C^A$ and $h, h^s \notin IN_C^N$ or $h, h^s \in IN_C^N$ and $h, h^s \notin IN_C^A$.

A basic handle assignment can be *completed*, i.e., turned into a complete handle assignment, by an obvious update of the IN_C ’s.

What the definition does not state yet is that IN_C ’s and the OUT_C ’s should be compatible, in the sense that the handles in IN_C^A and IN_C^N being active should not prevent the out-handles in OUT_C ’s from being true/false as required. Consider for instance the following extended cycle OC , which is meant to be a fragment of a wider program:

— OC

$a \leftarrow not\ b, f$

$b \leftarrow not\ c$

$c \leftarrow not\ a$

$b \leftarrow not\ e$

where $H_{OC} = \{(not\ e : OR : b), (f : AND : a)\}$.

Let us assume that $Out_handles(OC) = \{a, b\}$: this means, we assume that these are the atoms involved in OC that occur in the handles of some other cycle. Now take a handle assignment with the following components. $IN_{OC}^A = \{(f : AND : a)\}$ which means that we assume this handle to be active, i.e., we assume f to be false. $IN_{OC}^N = \{(not\ e : OR : b)\}$, which means that we assume this handle to be not active, i.e., we assume $not\ e$ to be false. Finally, $OUT_{OC}^+ = \{b\}$, and $OUT_{OC}^- = \{c\}$. For this handle assignment, the requirements about the out-handles are not compatible with the assumptions on the incoming handles. In fact, if f is assumed to be false, then a is concluded false, and consequently c is concluded true and b false (since moreover the OR handle $not\ e$ of b is in IN_{OC}^N , and thus is assumed to be not active). Notice that even with $IN_{OC}^N = \emptyset$ (i.e., with no knowledge about handle $(not\ e : OR : b)$) still with the given assumptions about the incoming handles we cannot assume to meet the requirements for OUT_{OC}^+ and OUT_{OC}^- . Instead,

with the same IN_{OC} 's, and with $OUT_{OC}^+ = \{c\}$ and $OUT_{OC}^- = \emptyset$ we obtain an handle assignment where the assumptions are compatible with the requirements. Notice also that $OUT_{OC}^- = \emptyset$ does not mean that no out-handle is *allowed* to be false, rather it means that no out-handle is *required* to be false. Then, provided that the requirements in OUT_{OC}^+ and OUT_{OC}^- are met, the remaining out-handles can take any truth value. Notice finally that if we let $IN_{OC}^A = IN_{OC}^N = \emptyset$, then the extended cycle is inconsistent.

Clearly, a definition of the IN_{OC} 's that makes the corresponding program fragment $C + Aux(C) + ActivationAt_C(IN_C^A)$ inconsistent is useless for obtaining stable models of the overall program. In fact, we are interested in handle assignments where the IN_{OC} 's correspond to an assumption on the incoming handles (and thus on $X_C = ActivationAt_C(IN_C^A)$) such that: the resulting program fragment $C + Aux(C) + X_C$ is consistent, and the requirements established in OUT_{OC}^+ and OUT_{OC}^- are met. This means that in some partial stable model of the program fragment all atoms in OUT_{OC}^+ are deemed true, and all atoms in OUT_{OC}^- are deemed false.

This is formalized in the following:

Definition 23

A handle assignment $HA = \langle IN_C^A, IN_C^N, OUT_C^+, OUT_C^- \rangle$ to a cycle C is *admissible* if and only if the program $C + Aux(C) + ActivationAt_C(IN_C^A)$ is consistent, and for some stable model $S^{IN_C^A}$ of this program, $OUT_C^+ \subseteq S^{IN_C^A}$ and $OUT_C^- \cap S^{IN_C^A} = \emptyset$. We say that $S^{IN_C^A}$ *corresponds* to HA .

According to Definition 14, each stable model $S^{IN_C^A}$ is a partial stable model of Π relative to C , that can be used for building a stable model of the whole program. At least some of these partial stable models *correspond* to the given handle assignment, in the sense that they are consistent with the choice of active handles that the assignment represents.

Proposition 1

A non-effective handle assignment cannot be admissible for an odd cycle.

Proof

A non-effective handle assignment provides an empty set of active handles to the cycle, which then is either unconstrained (no effective handle assignment exists because there are no handles) or is actually unconstrained (no handle is made active by this assignment). An unconstrained odd cycle is inconsistent. By Theorem 1, an actually unconstrained odd cycle is inconsistent as well, since it has no active handle. Then, by Definition 23 the assignment is not admissible. $\square$

and also that:

Proposition 2

A non-effective handle assignment is admissible for an even cycle C if and only if either $OUT_C^+ \subseteq Even_atoms(C)$ or $OUT_C^+ \subseteq Odd_atoms(C)$.

Proof

If cycle C is even, and it is either unconstrained or actually unconstrained, then the program fragment $C + Aux(C) + ActivationAt_C(IN_C^A)$ has two stable models, S_1 coinciding with $Even_atoms(C)$, and S_2 coinciding with $Odd_atoms(C)$. This because the handles either do not exist or are not active, and thus do not affect the stable models. By Definition 23 the assignment is effective if and only if $OUT_C^+ \subseteq S_1$ or $OUT_C^+ \subseteq S_2$. $\square$

Observe that whenever a handle assignment is effective the corresponding program fragment is locally stratified, and thus, according to (Przymusińska and Przymusiński 1990), has a unique stable model that coincides with its well-founded model. It may also be observed that a trivial handle assignment, which does not state requirements on the out-handles, is always admissible for an even cycle, and it is admissible for an odd cycle only if it is effective (otherwise as seen before the cycle is inconsistent).

The admissibility of a non-trivial effective handle assignment for cycle C can be checked syntactically, by means of the criterion that we state below. The advantage of this check is that it does not require to compute the well-founded model of $C + Aux(C) + ActivationAt_C(IN_C^A)$, but it just looks at the rules of C . Although the syntactic formulation may seem somewhat complex, it simply states in which cases an atom in OUT_C^+ , which is required to be concluded true w.r.t. the given handle assignment (or, conversely, an atom in OUT_C^- which is required to be concluded false), is actually allowed to take the specified truth value without raising inconsistencies. Notice that OUT_C^+ and OUT_C^- must be mutually coherent, in the sense that truth of an atom in OUT_C^+ cannot rely on truth of an atom in OUT_C^- (that instead is required to be concluded false), and vice versa.

Proposition 3

A non-trivial effective handle assignment $\langle IN_C^A, IN_C^N, OUT_C^+, OUT_C^- \rangle$ to cycle C is admissible if and only if for every $\lambda_i \in OUT_C^+$ the following condition (1) holds, and for every $\lambda_k \in OUT_C^-$ the following condition (2) holds.

1. Condition 1.

- (a) Either there exists OR handle h_o for λ_i , $h_o \in IN_C^A$ or
- (b) for every AND handle h_a for λ_i , $h_a \in IN_C^N$ and $\lambda_{i+1} \notin OUT_C^+$, and condition (2) holds for λ_{i+1} .

2. Condition 2.

- (a) For every OR handle h_o for λ , $h_o \in IN_C^N$, and
- (b) either there exists AND handle h_a for λ such that $h_a \in IN_C^A$, or $\lambda_{k+1} \notin OUT_C^-$, and condition (1) holds for λ_{k+1} .

Proof

Let us first notice that the set of rules with head λ_i in $C + Aux(C) + ActivationAt_C(IN_C^A)$ consists of rule $\lambda_i \leftarrow not \lambda_{i+1}, \Delta_i$ in C , and possibly, of one or more rules in $Aux(C)$. In fact, by the definition of canonical program, atoms in IN_C^A do not occur in C , and thus λ_i cannot belong to $ActivationAt_C(IN_C^A)$.

Consider an atom $\lambda_i \in OUT_C^+$, that we want to be concluded true in the partial stable model of C which corresponds to the given handle assignment. For λ_i to be concluded true, there must be a rule whose conditions are true w.r.t the handle assignment.

One possibility, formalized in Condition 1.(a), is that there exists an OR handle h_o for λ_i , $h_o \in IN_C^A$. That is, there is an auxiliary rule with head λ_i , and condition true w.r.t. the handle assignment.

Otherwise, as formalized in Condition 1.(b) we have to consider the rule of cycle C :

$$\lambda_i \leftarrow \text{not } \lambda_{i+1}, \Delta_i$$

and check that all the conditions are guaranteed to be true by the handle assignment. First of all it must be $(\Delta_k : AND : \lambda_k) \in IN_C^N$ i.e., in the given handle assignment the AND handle referring to λ_i must be not active, because an active AND handle would make the head of the rule false. Second, $\text{not } \lambda_{i+1}$ must be true: this on the one hand requires $\lambda_{i+1} \notin OUT_C^+$, that would be a contradiction; on the other hand, requires λ_{i+1} to be concluded false. To this aim, condition (2), discussed below, must hold for λ_{i+1} .

Consider now an atom $\lambda_k \in OC_C^-$, that we want to be false the partial stable model of C , which corresponds to the given assignment: there must *not* be a rule for λ_k whose conditions all true w.r.t. the given assignment.

First, as formalized in Condition 2.(a), we must have any OR handle h_o for λ_k in IN_C^N . Otherwise, λ_k would be necessarily concluded true, being the head of an auxiliary rule with a true body.

Second, as formalized in Condition 2.(b), we also have to consider the rule of cycle C

$$\lambda_k \leftarrow \text{not } \lambda_{k+1}, \Delta_k$$

and check that one of its two conditions is false w.r.t. the handle assignment. A first case is that $(\Delta_k : AND : \lambda_k) \in IN_C^A$, which means that the AND handle referring to λ_k is supposed to be active, i.e., false. Otherwise, $\text{not } \lambda_{k+1}$ must be false, i.e., λ_{k+1} must be true. To this aim, provided that $\lambda_{k+1} \notin OUT_C^-$ (that would be a contradiction), condition (1) must hold for λ_{k+1} . $\square$

The fact that Conditions 1 and 2 refer to each other is not surprising, since they are to be applied to cycles. Consider for instance the following cycle:

$$\begin{aligned} e &\leftarrow \text{not } f \\ f &\leftarrow \text{not } g \\ g &\leftarrow \text{not } e \\ g &\leftarrow h \end{aligned}$$

The handle assignment $\langle \{(h : OR)\}, \emptyset, \{g\}, \emptyset \rangle$ is admissible by Proposition 3, since, according to Condition 1.(a), there exists an auxiliary rule with head g and body in IN_C^A . Also $\langle \{(h : OR)\}, \emptyset, \{g, e\}, \emptyset \rangle$ is admissible, because: g is as above; there is no OR handle for e , thus Condition 1.(a) cannot be applied, but, considering rule $e \leftarrow \text{not } f$ (Condition 1.(b)), it is easy to see that Condition 2 holds of f , since there is no OR handle for f , and we have just shown that Condition 1 holds of g . Instead, $\langle \{(h : OR)\}, \emptyset, \{g\}, \{e\} \rangle$ is not admissible, because Condition 2 does not hold of e .

It is important to notice that it is possible to determine admissible handle assignments from a partially specified one. An obvious way of doing that is: guessing the missing sets, and checking whether the resulting handle assignment is admissible. There is however a much easier way by exploiting the definitions.

Namely, for given IN_C 's it is easy to find the maximal values for OUT_C^+ and OUT_C^- that form an admissible handle assignment. If IN_C^A is empty, then they correspond to the stable models (if any) of the cycle taken by itself (without the auxiliary rules, since an empty IN_C^A means that no OR handle is active). If IN_C^A is not empty, then by asserting the atoms in $ActivationAt_C(IN_C^A)$ as facts one computes the (unique) stable model of the extended cycle, and thus the maximal values for OUT_C^+ and OUT_C^- . These maximal values are determined by assuming all handles not belonging to the IN_C 's to be not active.

Vice versa, given OUT_C^+ and OUT_C^- , and unknown or partially defined IN_C 's, the conditions stated in Proposition 3 can be used for determining the subsets of H_C (incoming handles) that form admissible handle assignments.

Consider for instance the extended cycle:

$$\begin{aligned} e &\leftarrow \text{not } f, \text{not } r \\ f &\leftarrow \text{not } g \\ g &\leftarrow \text{not } e \\ g &\leftarrow v \\ g &\leftarrow h \\ e &\leftarrow s \\ e &\leftarrow \text{not } h \end{aligned}$$

and let $OUT_C^+ = \{g\}$ and $OUT_C^- = \emptyset$. Then, for forming an admissible handle assignment we have three possibilities.

First, by Condition 1.(a) of Proposition 3, we can exploit the auxiliary rule $g \leftarrow v$, i.e. the handle $(v : OR)$, and let $IN_C^{A1} = \{(v : OR)\}$, and $IN_C^{N1} = \emptyset$.

Second, again by Condition 1.(a) of Proposition 3, we can exploit the other auxiliary rule $g \leftarrow h$, i.e. the handle $(h : OR)$, and let $IN_C^{A2} = \{(h : OR)\}$. This implies to insert into IN_C^{N2} the contrary and opposite handles, since they both occur in C , i.e. $IN_C^{N2} = \{(h : AND), (\text{not } h : OR)\}$.

Third, we can exploit condition 1.(b), and consider rule with head g in the cycle, i.e. $g \leftarrow \text{not } e$, and verify Condition 2 for e , that must be false. For checking Condition 2.(a), we have to consider both the OR handles for e , i.e. handle $(\text{not } h : OR)$ and handle $(s : OR)$, that must be included in IN_C^{N2} , i.e., $IN_C^{N3} = \{(\text{not } h : OR), (s : OR)\}$. For checking Condition 2.(b) we have to consider rule $e \leftarrow \text{not } f, \text{not } r$. Since we want g to be true, this implies f to be false, which means that for getting e false as well, we have to add its AND handle $(\text{not } r : AND)$ to IN_C^{N3} . I.e., finally we get $IN_C^{N3} = \{(\text{not } h : OR), (s : OR), (\text{not } r : AND)\}$. This leads to add the opposite and contrary handles which occur in C to IN_C^{A2} , thus letting: $IN_C^{A3} = \{(h : OR)\}$.

We may notice that $IN_C^{A2} = IN_C^{A3}$ but $IN_C^{N2} \subseteq IN_C^{N3}$. Both choices form an admissible handle assignment, although the first one is more restricted. It turns out in fact that, in the above cycle, for building handle assignments where $OUT_C^+ = \{g\}$ and $OUT_C^- = \emptyset$, the

handle (*not* $r : AND$) is actually irrelevant. This explains why the definition of handle assignment does not enforce one to set *all* the handles of the cycle as active/not active. We can introduce the following definition:

Definition 24

An admissible handle assignment $\langle IN_C^A, IN_C^N, OUT_C^+, OUT_C^- \rangle$ is *minimal* if there is no other sets $IN_C^{A'} \subset IN_C^A$ and $IN_C^{N'} \subset IN_C^N$ such that $\langle IN_C^{A'}, IN_C^{N'}, OUT_C^+, OUT_C^- \rangle$ is still admissible.

As we have just seen, there can be alternative minimal sets of incoming active handles for the same out-handles. However, there may also be the case there is none. There is for instance no possibility for $OUT_C^+ = \{g, f\}$, i.e., no choice for the IN_C 's can produce a partial stable model where both g and f are true.

8 Cycle Graph and support sets

In previous sections we have proved that a stable model of a program can be obtained as the union of a compatible collection of partial stable models of the composing cycles. Partial stable models of a cycle are obtained by considering the corresponding extended cycle as a program, and making assumptions about its handles. We have discussed how to study consistency of extended cycles by means of the notion of handle assignment. In this section we introduce the Cycle Graph of a program, that represent cycles and handles. In the rest of the paper we show that the concepts and principles that we have previously introduced allow us to define syntactic conditions for consistency on the Cycle Graph.

Definition 25

Given program Π , the *Cycle Graph* CG_Π , is a directed graph defined as follows:

- **Vertices.** One vertex for each of cycles $C_1, \dots, C_w$ that occur in Π . Vertices corresponding to even cycles are labeled as EC_i 's while those corresponding to odd cycles are labeled as OC_j 's.
- **Edges.** An edge (C_j, C_i) marked with $(\Delta : K : \lambda)$ for each handle $(\Delta : K : \lambda) \in H_{C_i}$ of cycle C_i , that comes from C_j .

Each marked edge will be denoted by $(C_j, C_i | \Delta : K : \lambda)$, where either $(C_j$ or C_i or $\lambda)$ will be omitted whenever they are clear from the context, and we may write for short $(C_j, C_i | h)$, h standing for a handle that is either clear from the context or does not matter in that point. An edge on the CG connects the cycle a handle comes from to the cycle to which the handle belongs.

Take for instance the following program π_0 .

$\neg OC_1$
 $p \leftarrow \text{not } p, \text{not } c$
 $\neg Aux(OC_1)$
 $p \leftarrow \text{not } b$
 $\neg OC_2$
 $q \leftarrow \text{not } q$
 $\neg Aux(OC_2)$
 $q \leftarrow a$
 $q \leftarrow \text{not } e$
 $\neg OC_3$
 $r \leftarrow \text{not } r, \text{not } e$
 $\neg EC_1$
 $c \leftarrow \text{not } d$
 $d \leftarrow \text{not } c$
 $\neg EC_2$
 $a \leftarrow \text{not } b$
 $b \leftarrow \text{not } a$
 $\neg EC_3$
 $e \leftarrow \text{not } f$
 $f \leftarrow \text{not } e$

Its cycle graph CG_{π_0} is shown in Figure 1.




Figure 1. The Cycle Graph of π_0 .

The Cycle Graph of a program directly represents cycles, that correspond to its vertices. It also indirectly represents extended cycles, since its edges are marked by the handles. Paths on the Cycle Graph graph represent direct or indirect connections between cycles through the handles. In order to relate admissible handle assignments for the cycles of Π to subgraphs of its cycle graph CG_{Π} we introduce the following definitions.

Definition 26

Given program Π , let a *CG support set* be a pair $S = \langle ACT^+, ACT^- \rangle$ of subsets of the

handles marking the edges of CG_Π (represented in the form $(\Delta : K)$ with $K = AND/OR$) such that the following conditions hold:

- (i) $ACT^+ \cap ACT^- = \emptyset$.
- (ii) neither ACT^+ nor ACT^- contain a pair of either opposite or contrary handles.
- (iii) if two opposite handles h and h^- both occur on the CG , then ACT^+ contains handle h if and only if ACT^- contains its opposite handle h^- .
- (iv) if two contrary handles h and h^n both occur on the CG , then ACT^+ contains handle h if and only if ACT^- contains its contrary handle h^n .
- (v) if two sibling handles h and h^s both occur on the CG , then either $h, h^s \in ACT^+$ and $h, h^s \notin ACT^-$, or vice versa $h, h^s \in ACT^-$ and $h, h^s \notin ACT^+$.

Given S , we will indicate its two components with $ACT^+(S)$ and $ACT^-(S)$. For the sake of readability we introduce some simplifying assumptions.

- Given handle $h = (\Delta : K : \lambda)$, by $ACT^+(S) \cup \{h\}$ (resp. $ACT^-(S) \cup \{h\}$) we mean $ACT^+(S) \cup \{(\Delta : K)\}$ (resp. $ACT^-(S) \cup \{(\Delta : K)\}$).
- Given handle $h \in ACT^+(S)$ (resp. $h \in ACT^-(S)$) of the form $(\Delta : K)$, by $IN_C^A \cup \{h\}$ (resp. $IN_C^N \cup \{h\}$) we mean: to identify the set $H = \{(\Delta : K : \lambda) \in H_C\}$ and perform $IN_C^A \cup H$ (resp. $IN_C^N \cup H$).
- By $H_C \cap ACT^+(S)$ (resp. $H_C \cap ACT^-(S)$) we mean $\{(\Delta : K : \lambda) \in H_C \mid (\Delta : K) \in ACT^+(S)\}$ (resp. $\{(\Delta : K : \lambda) \in H_C \mid (\Delta : K) \in ACT^-(S)\}$).

As stated in Theorem 1, we have to restrict the attention on CG support sets including at least one active handle for each odd cycle. Then, according to Theorem 2, we have to check that the assumptions on the handles are mutually coherent, and are sufficient for ensuring consistency.

Definition 27

A CG support set S is *potentially adequate* if for every odd cycle C in Π there exists a handle $h \in H_C$ such that $h \in ACT^+(S)$.

A CG support set S induces a set of handle assignments, one for each of the cycles $\{C_1, \dots, C_w\}$ occurring in Π .

The induced assignments are obtained on the basis of the following observations:

- Each handle in $h \in ACT^+(S)$ is supposed to be active, and therefore it must be active for each cycle C_i such that $h \in H_{C_i}$.
- Each handle in $h \in ACT^-(S)$ is supposed to be not active, and therefore it must be not active for each of cycle C_j such that $h \in H_{C_j}$.
- If a handle h in S requires, in order to be active/not active, an atom β to be false, then it must be concluded false *in all the extended cycles of the program* h comes from.
- If a handle h in S requires, in order to be active/not active, an atom β to be true, then it must be concluded true *in all the extended cycles of the program* h comes from. This point deserves some comment, since one usually assumes that it suffices to conclude β true *somewhere* in the program. Consider however that any rule $\beta \leftarrow \text{Body}$ that allows β to be concluded true in some cycle is an auxiliary rule to all the other cycles

β is involved into. This is why β is concluded true *everywhere it occurs*. This is the mechanism for selecting partial stable models of the cycles that agree on shared atoms, in order to assemble stable models of the overall program.

Definition 28

Let $S = \langle ACT^+, ACT^- \rangle$ be a GG support set which is potentially adequate. For each cycle C_k occurring in Π , $k \leq w$, the (possibly empty) handle assignment induced by this set is determined as follows.

1. Let $IN_{C_k}^A$ be $H_{C_k} \cap ACT^+(S)$.
2. Let $IN_{C_k}^N$ be $H_{C_k} \cap ACT^-(S)$.
3. Let $OUT_{C_k}^+$ be the (possibly empty) set of all atoms $\beta \in Out_handles(C_k)$ such that there is a handle $h \in ACT^+(S)$ either of the form $(\beta : OR)$ or $(not \beta : AND)$.
4. Let $OUT_{C_k}^-$ be the (possibly empty) set of all atoms $\alpha \in Out_handles(C_k)$ such that there is a handle $h \in ACT^-(S)$ either of the form $(\alpha : AND)$ or $(not \alpha : OR)$.
5. Verify that $OUT_{C_k}^- \cap OUT_{C_k}^+ = \emptyset$.

If this is the case for each C_k , then S actually induces a set of handle assignments, and is called *coherent*. Otherwise, S does not induce a set of handle assignments, and is called *incoherent*.

The above definition does not guarantee that the assignments induced by a coherent support set are admissible, that the same atom is not required to be both true and false in the assignments of different cycles, and that the incoming handles of a cycle being supposed to be active/not active corresponds to a suitable setting of the out-handles of the cycles they come from. Consider for instance cycle C_i which has an incoming handle, e.g. $h = (\beta : OR : \lambda)$, in $IN_{C_i}^A$: h is supposed to be active, which in turn means that β must be concluded true elsewhere in the program; then, for all cycles C_j where β is involved into, we must have $\beta \in OUT_{C_j}^+$, in order to fulfill the requirement. Of course, we have to consider both IN_C^A and IN_N^C , and both the AND and the OR handles.

The following definition formalizes these more strict requirements.

Definition 29

A coherent CG support set S is *adequate* (w.r.t. not adequate) if for the induced handle assignments the following conditions hold:

1. they are admissible;
2. for each two cycles C_i, C_j in Π , $OUT_{C_i}^+ \cap OUT_{C_j}^- = \emptyset$.
3. For every C_i in Π , for every handle $h \in IN_{C_i}^A$ of the form either $(\beta : OR : \lambda)$ or $(not \beta : AND : \lambda)$, and for every handle $h \in IN_{C_i}^N$ of the form either $(\beta : AND : \lambda)$ or $(not \beta : OR : \lambda)$, for every other cycle C_j in Π , $i \neq j$, such that $\beta \in Out_handles(C_j)$, we have $\beta \in OUT_{C_j}^+$.

4. For every C_i in Π , for every handle $h \in IN_{C_k}^A$ of the form either $(not \beta : OR : \lambda)$ or $(\beta : AND : \lambda)$, and for every handle $h \in IN_{C_k}^N$ of the form either $(not \beta : AND : \lambda)$ or $(\beta : OR : \lambda)$, for every other cycle C_j in Π , $i \neq j$, such that $\beta \in Out_handles(C_j)$, we have $\beta \in OUT_{C_j}^-$.

9 Checking consistency on the Cycle Graph: Main Result

We are now able to state the main result of the paper, which gives us a necessary and sufficient syntactic condition for consistency based on the Cycle Graph of the program.

Theorem 3

A program Π has stable models if and only if there exists an adequate CG support set S for Π .

Proof

$\Leftarrow$

On the basis of S we can obtain the corresponding induced handle assignments, that will be admissible by the hypothesis that the S is adequate.

From S we can obtain a *global handle assignment* $HA = \langle T_{HA}; F_{HA} \rangle$ as follows.

$$T_{HA} = \{\alpha \mid (\alpha : OR) \in ACT^+(S) \vee (not \alpha : AND) \in ACT^+(S) \vee (not \alpha : OR) \in ACT^-(S) \vee (\alpha : AND) \in ACT^-(S)\}$$

$$F_{HA} = \{\alpha \mid (\alpha : AND) \in ACT^+(S) \vee (not \alpha : OR) \in ACT^+(S) \vee (\alpha : OR) \in ACT^-(S) \vee (not \alpha : AND) \in ACT^-(S)\}$$

Since the S is adequate, then by Definition 29 we have that (i) for each cycle C_i in Π , S induces an admissible handle assignment, that (ii) HA is consistent (i.e. $T_{HA} \cap F_{HA} = \emptyset$), that (iii) $\forall \alpha \in T_{HA}$, α is concluded true in every cycle C_i it is involved into since $\alpha \in OUT_{C_i}^+$, and the handle assignment induced by S to C_i is admissible, and that (iv) $\forall \alpha \in F_{HA}$, α is concluded false in every cycle C_j it is involved into, since $\alpha \in OUT_{C_j}^+$, and the handle assignment induced by S to C_j is admissible.

On the basis of HA , for each cycle C_i in Π we can build a correspondent independent program $C_i + Aux(C_i) + X_i$, where we let $X_i = Atoms(H_{C_i}) \cap T_{HA}$, and $X_i^- = Atoms(H_{C_i}) \cap F_{HA}$. This independent program has a stable model S_i by construction, since the handle assignment induced by S to C_i is admissible (point (i) above). This stable model is (by Definition 14) a partial stable model for Π relative to C_i , with base $\langle X_i, X_i^- \rangle$. Taken one S_i for each C_i in Π , in the terminology of Definition 15 the S_i 's constitute a compatible set of partial stable models because, according to Definition 15: (1) for any other cycle C_j , $X_i \cap X_j^- = \emptyset$, since $T_{HA} \cap F_{HA} = \emptyset$ by point (ii) above; (2) $\forall A \in X_i$, A is concluded true in some other cycle, by point (iii) above; (3) $\forall A \in X_i^-$, A is concluded false all the other cycles, by point (iv) above. Then, by Theorem 2, Π has stable models.

$\Rightarrow$

If Π has a stable model M , then by Theorem 2 we can decompose M into a compatible set

of partial stable models, one partial stable model S_i , with base $\langle X_i, X_i^- \rangle$, for each C_i in Π . Since $X_i, X_i^- \subseteq \text{Atoms}(H_{C_i})$, they correspond to sets $IN_{C_i}^A$ and $IN_{C_i}^N$ of handles of C_i that are made active/not active by this base. By point (1) of Definition 15, for every two cycles C_i, C_j $X_i \cap X_j^- = \emptyset$, and then $IN_{C_i}^A \cap IN_{C_j}^N = \emptyset$. If we let S such that $ACT^+(S) = \bigcup_{i \leq w} IN_{C_i}^A$ and $ACT^-(S) = \bigcup_{i \leq w} IN_{C_i}^N$, we have $ACT^+(S) \cap ACT^-(S) = \emptyset$ and, if there are either opposite or contrary handles, they will not be in the same set. Then, S is a CG support set, and is potentially adequate by construction, because it has been built from the IN^A 's and IN^N 's of the cycles (point 1-2 of Definition 27), and because the S_i 's agree on shared atoms, having been obtained by decomposing a stable model (points 3-5 of Definition 27). For the same reasons, S is also adequate.

□

Checking the condition stated in Theorem 3 does not imply finding the stable models of the program. However, in the proof of the *only-if* part, a way of determining the stable models can be actually outlined, and is summarized below.

Corollary 1

Assume that the condition stated in Theorem 3 holds for program Π . Then, the stable models of Π can be determined as follows.

1. Given the handle assignments induced by S , build T_{HA} .
2. For each cycle C_i in Π , let $X_i = \text{Atoms}(H_{C_i}) \cap T_{HA}$, build the corresponding extended cycle $C_i + \text{Aux}(C_i) + X_i$, and find its partial stable models.
3. Assemble each stable model of Π as the union of one partial stable model for each cycle.

Consider the following collection of cycles.

— OC_0

$p \leftarrow \text{not } s, \text{not } c$

$s \leftarrow \text{not } t$

$t \leftarrow \text{not } p$

— OC_1

$p \leftarrow \text{not } s, \text{not } c$

$s \leftarrow \text{not } t$

$t \leftarrow \text{not } p$

$s \leftarrow a$

— OC_2

$q \leftarrow \text{not } q$

$q \leftarrow \text{not } e$

— OC_3

$r \leftarrow \text{not } r, \text{not } e$

— EC_1
 $a \leftarrow \text{not } c$
 $c \leftarrow \text{not } a$

— EC_2
 $a \leftarrow \text{not } b$
 $b \leftarrow \text{not } a$

— EC_3
 $e \leftarrow \text{not } f$
 $f \leftarrow \text{not } e$

— OC'_2
 $q \leftarrow \text{not } q$
 $q \leftarrow \text{not } e$
 $q \leftarrow a$

Let $\pi_1 = OC_0 \cup EC_1$, $\pi_2 = OC_1 \cup EC_1 \cup EC_2$, $\pi_3 = OC_1 \cup EC_1 \cup EC_2 \cup OC_2 \cup OC_3$, and $\pi_4 = OC_1 \cup EC_1 \cup EC_2 \cup OC'_2 \cup OC_3$. The Cycle Graphs of these programs are shown in Figure 9, Figures 2, 3 and 4 respectively.






Figure 2. The Cycle Graph of π_1 .

Let's denote $C + Aux(C)$ by C^e .

For π_1 , we have $T_{HA} = \{c\}$. The partial stable model of $EC_1^e \cup c$ is $\{c\}$; the partial stable model of $OC_0^e \cup c$ is $\{c, t\}$. Then, a stable model of the overall program is (as it is easy to verify) $\{c, t\}$.


For π_2 , we have two possibilities. In the first one, $T_{HA} = \{c\}$. The partial stable model of $EC_1^e \cup c$ is $\{c\}$; the partial stable model of $EC_2^e \cup c$ is $\{c, b\}$; the partial stable model of $OC_1^e \cup c$ is $\{c, t\}$. Then, a stable model of the overall program is (as it is easy to verify) $\{c, b, t\}$.

Figure 3. The Cycle Graph of π_2 .Figure 4. The Cycle Graph of π_3 .

In the second one, $T_{HA} = \{a\}$. The partial stable model of $EC_1^e \cup a$ is $\{a\}$; the partial stable model of $EC_2^e \cup a$ is $\{a\}$; the partial stable model of $OC_1^e \cup a$ is $\{a, s, t\}$. Then, a stable model of the overall program is (as it is easy to verify) $\{a, s, t\}$.

For π_3 the situation is hopeless, since the only incoming handles to OC_2 and OC_3 are opposite handles, that cannot be both active.

For π_4 , we have $T_{HA} = \{a, e\}$. The partial stable model of $EC_1^e \cup a$ is $\{a\}$; the partial stable model of $EC_2^e \cup a$ is $\{a\}$; the partial stable model of $OC_1^e \cup a$ is $\{a, s, t\}$; the partial stable model of $OC_2^e \cup a$ is $\{a, q\}$; the partial stable model of $EC_3^e \cup e$ is $\{e\}$; the partial stable model of $OC_3^e \cup e$ is $\{e\}$. Then, a stable model of the overall program is (as it is easy to verify) $\{a, s, t, q, e\}$.

Figure 5. The Cycle Graph of π_4 .

Therefore, it is useful to define a procedure for identifying adequate support sets of a program on its Cycle Graph.

10 Identifying adequate support sets on the Cycle Graph

The above definitions allow us to define a procedure for trying to *find* adequate support sets starting from the odd cycles, and following the dependencies on the CG.

Definition 30 (Procedure PACG for finding adequate CG support sets for program Π)

1. Let initially $S = \langle \emptyset; \emptyset \rangle$.
2. For each cycle C_k occurring in Π , $k \leq w$, let initially $HA_{C_k} = \langle \emptyset, \emptyset, \emptyset, \emptyset \rangle$
3. For each odd cycle OC in Π do:
 - (a) Choose $h \in H_{OC}$. If $H_{OC} = \emptyset$, than FAIL.
 - (b) For chosen h :
 - i do $ACT^+(S) := ACT^+(S) \cup \{h\}$;
 - ii if h^s occurs in the CG, do $ACT^+(S) := ACT^+(S) \cup \{h^s\}$;
 - iii if h^- occurs in the CG, do $ACT^-(S) := ACT^-(S) \cup \{h^-\}$;
 - iv if h^n occurs in the CG, do $ACT^-(S) := ACT^-(S) \cup \{h^n\}$.
 - v For each cycle C_k in Π such that $h \in H_{C_k}$:
 - A do $IN_{C_k}^A := IN_{C_k}^A \cup h$;
 - B if h^s occurs in H_{C_j} for some cycle C_j (where possibly $j = k$), do $IN_{C_j}^A := IN_{C_j}^A \cup \{h^s\}$;
 - C if h^- occurs in H_{C_j} for some cycle C_j (where possibly $j = k$), do $IN_{C_j}^N := IN_{C_j}^N \cup \{h^-\}$;

- D if h^n occurs in H_{C_j} for some cycle C_j (where possibly $j = k$), do $IN_{C_j}^N := IN_{C_j}^N \cup \{h^n\}$.
- vi If h is either of the form $(\beta : OR)$ or $(not \beta : AND)$, for each cycle C_k in Π where $\beta \in Out_handles(C_k)$, do:
 $OUT_{C_k}^+ := OUT_{C_k}^+ \cup \{\beta\}$;
- vii If h is either of the form $(not \beta : OR)$ or $(\beta : AND)$, for each cycle C_k in Π where $\beta \in Out_handles(C_k)$, do:
 $OUT_{C_k}^- := OUT_{C_k}^- \cup \{\beta\}$

4. REPEAT

- (a) Verify that $ACT^+(S) \cap ACT^-(S) = \emptyset$. If not, FAIL.
- (b) Verify that neither ACT^+ nor ACT^- contain a pair of either opposite or contrary handles. If not, FAIL.
- (c) For each cycle C_k in Π such that $OUT_{C_k}^+ \neq \emptyset$ or $OUT_{C_k}^- \neq \emptyset$:
 - i Verify that $OUT_{C_k}^+ \cap OUT_{C_k}^- = \emptyset$. If not, FAIL.
 - ii Update (if needed) $IN_{C_k}^A$ and $IN_{C_k}^N$ w.r.t. $OUT_{C_k}^+$ and $OUT_{C_k}^-$, and check that the resulting handle assignment is admissible. If not, then FAIL.
 - iii For each other cycle C_h in Π do: verify that $OUT_{C_k}^+ \cap OUT_{C_h}^- = \emptyset$, and that $OUT_{C_k}^- \cap OUT_{C_h}^+ = \emptyset$. If not, FAIL.
- (d) For each cycle C_k in Π , for each $h \in IN_{C_k}^A$:
 - i do $ACT^+(S) := ACT^+(S) \cup \{h\}$;
 - ii if h^s occurs in the CG, do $ACT^+(S) := ACT^+(S) \cup \{h^s\}$;
 - iii if h^- occurs in the CG, do $ACT^-(S) := ACT^-(S) \cup \{h^-\}$;
 - iv if h^n occurs in the CG, do $ACT^-(S) := ACT^-(S) \cup \{h^n\}$.
 - v For each cycle C_h in Π such that $h \in H_{C_h}$:
 - A do $IN_{C_h}^A := IN_{C_h}^A \cup h$;
 - B if h^s occurs in H_{C_j} for some cycle C_j (where possibly $j = h$), do $IN_{C_j}^A := IN_{C_j}^A \cup \{h^s\}$;
 - C if h^- occurs in H_{C_j} for some cycle C_j (where possibly $j = h$), do $IN_{C_j}^N := IN_{C_j}^N \cup \{h^-\}$;
 - D if h^n occurs in H_{C_j} for some cycle C_j (where possibly $j = h$), do $IN_{C_j}^N := IN_{C_j}^N \cup \{h^n\}$.
 - vi If h is either of the form $(\beta : OR)$ or $(not \beta : AND)$, for each cycle C_k in Π where $\beta \in Out_handles(C_k)$, do:
 $OUT_{C_k}^+ := OUT_{C_k}^+ \cup \{\beta\}$;
 - vii If h is either of the form $(not \beta : OR)$ or $(\beta : AND)$, for each cycle C_k in Π where $\beta \in Out_handles(C_k)$, do:
 $OUT_{C_k}^- := OUT_{C_k}^- \cup \{\beta\}$
- (e) For each cycle C_k in Π , for each $h \in IN_{C_k}^N$:
 - i do $ACT^-(S) := ACT^-(S) \cup \{h\}$;

- ii if h^s occurs in the CG, do $ACT^-(S) := ACT^-(S) \cup \{h^s\}$;
- iii if h^- occurs in the CG, do $ACT^+(S) := ACT^-(S) \cup \{h^-\}$;
- iv if h^n occurs in the CG, do $ACT^+(S) := ACT^+(S) \cup \{h^n\}$.
- v For each cycle C_h in Π such that $h \in H_{C_h}$:
 - A do $IN_{C_h}^N := IN_{C_h}^N \cup h$;
 - B if h^s occurs in H_{C_j} for some cycle C_j (where possibly $j = h$), do $IN_{C_j}^N := IN_{C_j}^N \cup \{h^s\}$;
 - C if h^- occurs in H_{C_j} for some cycle C_j (where possibly $j = h$), do $IN_{C_j}^A := IN_{C_j}^A \cup \{h^-\}$;
 - D if h^n occurs in H_{C_j} for some cycle C_j (where possibly $j = h$), do $IN_{C_j}^A := IN_{C_j}^A \cup \{h^n\}$.
- vi If h is either of the form $(\beta : OR)$ or $(not \beta : AND)$, for each cycle C_k in Π where $\beta \in Out_handles(C_k)$, do: $OUT_{C_k}^- := OUT_{C_k}^- \cup \{\beta\}$;
- vii If h is either of the form $(not \beta : OR)$ or $(\beta : AND)$, for each cycle C_k in Π where $\beta \in Out_handles(C_k)$, do: $OUT_{C_k}^+ := OUT_{C_k}^+ \cup \{\beta\}$

UNTIL no set is updated by the previous steps.

Proposition 4

Procedure PACG either fails, or returns an adequate CG support set.

Proof

Whenever it does not fail, PACG clearly produces a CG support set S . In fact: points (i-ii) of Definition 26 are verified by steps 4.(a-b) of PACG; points (iii-v) of Definition 26 are enforced after any update to S , namely by steps 3.b.(ii-iv), 4.d.(ii-iv) and 4.e.(ii-iv). The CG support set S produced by PACG is potentially adequate by construction, since in step 3.a a handle for each odd cycle is included. S is also adequate, since in fact: admissible handle assignments for all cycles in Π are incrementally built and verified in steps 4.c.(i-ii), thus fulfilling point 1. of Definition 29. Point 2 of Definition 29 is verified in step 4.c.(iii). Finally, points 3-4 of Definition 29 are enforced by steps 3.b.(vi-vii), 4.d.(vi-vii) and 4.e.(vi-vii), after each update to the IN_C 's of any cycle. $\square$

Let us reconsider collection of cycles given in previous sections, and programs π_1 , π_2 , π_3 and π_4 .

For $\pi_1 = OC_0 \cup EC_1$, the odd cycle OC_0 admits the unique potentially active handle $(not\ c : AND : p)$. Then, we let S_{π_1} be such that $ACT^+(S_{\pi_1}) = \{(not\ c : AND)\}$ and $ACT^-(S_{\pi_1}) = \emptyset$. The induced set of handle assignments are as follows.

For OC_0 : $IN_{OC_0}^A = \{(not\ c : AND)\}$, $OUT_{OC_0}^+ = OUT_{OC_0}^- = \emptyset$. This assignment is trivially admissible, since there is no requirement on the out-handles.

For EC_1 : $OUT_{EC_1}^+ = \{c\}$, $OUT_{EC_1}^- = \{\emptyset\}$. $IN_{EC_1} = \emptyset$, since EC_1 is unconstrained. It is easy to verify that this handle assignment is admissible, by letting $\lambda_1 = a$ and $\lambda_2 = c$,

where of course for c to be true a must be false. This handle assignment corresponds to selecting the partial stable model $\{c\}$ for EC_1 , while discarding the other partial stable model $\{a\}$.

Then, S_{π_1} as defined above is an adequate CG support set.

Consider program $\pi_2 = OC_1 \cup EC_1 \cup EC_2$. The situation here is complicated by the fact that EC_1 and EC_2 are not independent. In fact, rule $a \leftarrow \text{not } b$ of EC_2 is an auxiliary rule for EC_1 , and, vice versa, $a \leftarrow \text{not } c$ of EC_1 is an auxiliary rule for EC_2 . Then, here we have a cyclic connection between the even cycles. This is evident on the cycle graph of π_2 , reported in Figure 2.

The odd cycle OC_1 has two handles, of which at least one must be active. Let us first assume that $(\text{not } c : \text{AND} : p)$ is active. According to the PACG procedure, we try to assemble a CG support set S , by letting at first $ACT^+(S_{\pi_2}) = \{(\text{not } c : \text{AND})\}$ and $ACT^-(S_{\pi_2}) = \{(\text{not } c : \text{OR})\}$. In fact, since $\text{not } c$ is an incoming OR handle for a in EC_2 , when assuming $(\text{not } c : \text{AND})$ to be active, we also have to assume its opposite handle and its contrary handle to be not active.

Accordingly, we let $IN_{OC_1}^A = \{(\text{not } c : \text{AND})\}$ and $IN_{EC_2}^N = \{(\text{not } c : \text{OR})\}$. Now, we have to put $OUT_{OC_1}^- = \{p\}$ and $OUT_{EC_1}^+ = \{c\}$. To form an admissible handle assignment for EC_1 , this implies to let $IN_{EC_1}^N = \{(\text{not } b : \text{OR})\}$. Consequently, we have to update $ACT^-(S_{\pi_2})$ which becomes: $ACT^-(S_{\pi_2}) = \{(\text{not } c : \text{AND}), (\text{not } b : \text{OR})\}$. This leads to put $OUT_{EC_1}^+ = \{b\}$.

Further iteration of the procedure changes nothing, and thus the pair of sets $ACT^+(S_{\pi_2}) = \{(\text{not } c : \text{AND})\}$ and $ACT^-(S_{\pi_2}) = \{(\text{not } c : \text{OR}), (\text{not } b : \text{OR})\}$ form, as it is easy to verify, an adequate CG support set.

Notice that this kind of reasoning requires neither to find the stable models of the cycles, nor to consider every edge of the CG. In fact, we do not need to consider the second incoming handle of OC_1 .

Let us now make the alternative assumption, i.e. assume that $(a : \text{OR} : s)$ is active for OC_1 . This means at first $ACT^+(S_{\pi_2}) = \{(a : \text{OR})\}$ and $ACT^-(S_{\pi_2}) = \emptyset$, since $\text{not } a$ does not occur in handles of the CG. This implies $OUT_{EC_1}^+ = \{a\}$. Thus, there is no requirement on IN_{EC_2} for forming an admissible handle assignment, and then the procedure stops here.

For program $\pi_3 = OC_1 \cup EC_1 \cup EC_2 \cup OC_2 \cup OC_3$, as we have already seen the only incoming handles to OC_2 and OC_3 are opposite handles, that cannot be both active. For the other cycles, the situation is exactly as before.

Then, there is a subprogram which gives problems. We can fix these problems for instance by replacing OC_2 with OC'_2 , thus obtaining program π_4 (CG in Figure 4) where we can exploit handle $(a : \text{OR})$ for both OC_1 and OC'_2 . It is easy to verify that the CG support set S composed of $ACT^+(S_{\pi_4}) = \{(a : \text{OR}), (\text{not } e : \text{AND})\}$ and $ACT^-(S_{\pi_4}) = \{(\text{not } e : \text{OR})\}$ is adequate. The need to support OC'_2 rules out the possibility of supporting OC_1 by means of the handle $(\text{not } c : \text{AND} : p)$.

11 Usefulness of the results

We believe that our results can be useful in different directions: (i) Making consistency checking algorithms more efficient in the average case. (ii) Defining useful classes of programs which are consistent by construction. (iii) Checking properties of programs statically and dynamically, i.e., when modifications and updates affect the existence and the number of answer sets. (iv) Introducing component-based software engineering principles and methodologies for answer set programming: this by defining, over the *CG*, higher level graphs where vertices are components, consisting of bunches of cycles, and edges are the same of the *CG*, connecting components instead of single cycles.

11.1 Splitting consistency checking into stages

The approach and the result that we have presented here can lead to defining new algorithms for computing stable models. However, they can also be useful for improving existing algorithms.

We have identified and discussed in depth two aspects of consistency checking: (1) the odd cycles must be (either directly or indirectly) supported by the even cycles; (2) this support must be consistent, in the sense that no contrasting assumptions on the handles can be made.

Point (1) is related to the “coarse” structure of the program, and can be easily checked on the *CG*, so as to rule out a lot of inconsistent programs, thus leaving only the “potentially consistent” ones to be checked w.r.t. point (2). This is the aspect that might be potentially exploited by any approach to stable models computation.

Notice that a *CG* support set S determines a subgraph of the *CG*, which is composed of all the edges (and the corresponding end vertices) marked with the handles which occur in S .

Definition 31

Given the *CG* of program Π , and a *CG* support set S , an *adequate support subgraph* is a subgraph CG_S of the *CG*, composed of the edges marked by the handles belonging to $ACT^+(S)$ and $ACT^-(S)$, and of the vertices connected by these edges.

It is easy to see that, syntactically, CG_S is composed of a set of *handle paths*, that connect the odd cycles, through a chain of handles, to the even cycles (or to cyclic bunches of even cycles) that are able to support them. Each path may include more than one odd cycle, while each odd cycle must occur in at least one path.

Then, point 1 above may consist in checking whether a subgraph of the *CG* with this syntactic structure exists. Point 2, however performed, in essence must check whether the handles marking the subgraph constitute an adequate *CG* support set.

Staying within the approach of this paper, one may observe that the PACG procedure can easily be generalized for computing the stable models by performing the two steps in parallel. In fact, PACG actually tries to reconstruct the CG_S , starting from the odd cycles and going backwards through the *CG* edges to collect the handles that form the set S .

At each step however, the procedure updates the handle assignments of the cycles and performs the necessary checks to be sure to be assembling an adequate set S . The extension would consist in computing the stable models of the extended cycles instead of just the handle assignments, and perform the computation on the whole CG .

11.2 Defining classes of programs that are consistent by construction

Based on the CG it is possible to define syntactic restrictions that, with a slight loss of expressivity, may ensure the existence of stable models. Suitable restrictions might be enforced “on line” by an automated tool, while the program is being written. This can be made easier by limiting the number of handles each cycle may have.

The definition of classes of programs suitable for “interesting” applications is a topic of further research, but it can be useful to give some hints here.

In the literature, various sufficiency conditions have been defined (beyond stratification) for existence of stable models.

- Acyclic programs, by Apt and Bezem (Apt and Bezem 1991);
- Signed programs, by Turner (Turner 1994);
- Call-consistent programs, order consistent programs, and negative cycle free programs by Fages (Fages 1994).

For programs without classical negation, the classes of acyclic programs, signed programs, negative cycle free and order consistent programs are either included or coincide with the class of locally stratified programs (Przymusinska and Przymusinski 1990), and have a unique stable model that coincides with the well-founded model. In that case, their canonical counterpart is the empty program. Call-consistent programs do not contain odd cycles by definition.

We define below a new very simple class of programs that are guaranteed to have stable models, broader than the above ones since we admit odd cycles.

Definition 32

A program Π is called *tightly-even-bounded* if it is either call-consistent, or (if not) the following conditions hold: (i) every odd cycle has just one handle; (ii) this handle comes from an unconstrained even cycle; (iii) if there are two odd cycles whose handles come from the same even cycle, then two handles that originate in the same kind of node are of the same kind.

The above condition is clearly very easily and directly checked on the CG , and can be made clearly visible and understandable to a user, via a graphical interface. If you take any other existing graph representation, like e.g. the *EDG* (Brignoli et al. 1999) (Costantini 2001), that computes stable models as *graph colorings*, the check is of course possible, but is less easy and less direct.

It is easy to see that:

Theorem 4

Every tightly even-bounded program P has stable models.

Proof

Even cycles the handles come from are unconstrained, and conflicting handles are excluded by definition. Then, we can build an adequate CG support set by just assuming the incoming handles of the odd cycles to be active. $\square$

Simple as it is, this is a class wider than that of consistent programs, which is easy to understand by programmers, and is guaranteed to have stable models. Moreover, we can further enlarge this class by allowing a tightly even-bounded “core” program to have a “top”, i.e. a set of definitions that do not contain cycles, and possibly depend upon atoms of the “core” part. The resulting class of program has a *generate* part, consisting of the even cycles, a *test* part consisting of the odd cycles which “prune” stable models, and a *conclude* part that draws further consequences.

11.3 Checking Properties of Programs

By inspecting the structure of the CG it is possible in principle to detect whether a program is categorical, i.e. has a unique answer set, and to estimate the number of the answer sets. In past work, we have investigated the effects on the existence and the number of answer sets after modifications to the program. In particular, after asserting lemmas (Costantini et al. 1996), and if adding new rules to the program in the program development stage (Costantini et al. 2003). The above results can be reformulated, made uniform and extended by employing the CG for representing the program. Other static and dynamic program properties may be investigated.

11.4 Generalizing the CG to components/agents

A relevant topic is, in our opinion, that of defining software engineering principles for Answer Set Programming.

Here we propose to define a program development methodology for Answer Set Programming by defining, over the CG, higher level graphs where vertices are *components*, and edges are that of the CG, but connect components instead of single cycles. We give below a first informal description of what kind of methodology we actually mean.

Let a *component* C be a bunch of cycles. It can be developed on its own, or it can be identified on the CG of a larger program. Similarly to a cycle however, C is not meant to be an independent program, but rather it has incoming handles.

As we have seen, partial stable models of cycles are characterized by handle assignments. Analogously, a component will be characterized by a *component interface*

$$IN_C^A, IN_C^N, OUT_C^+, OUT_C^-$$

that is meant to be a specification of which values the incoming handles may take, either

in order to keep the component consistent, or in order to select some of its stable models. The out-handles provide the other components with a mean of establishing a connection with this one, i.e., they are true/false atoms that can make the incoming handles of other components active/not active as required.

Differently from cycles, components in general will not export *all* their active handles, but only those they want to make visible and available outside.

Based on the interface, it is possible to connect components, thus building a *Component Graph Comp_G*. On this new graph *Comp_CG*, one can either add new consistent components, or modify existing ones, and can check over the handle paths if there are problems for consistency, and how to fix them.

Referring to the previous example, in π_3 we have the component $OC_1 \cup EC_1 \cup EC_2$ which is consistent, and the component $OC_2 \cup OC_3 \cup EC_3$ which is instead inconsistent. Then, we have a *Comp_CG* with two unconnected vertices. We have shown how to fix the problem by adding a handle to OC_2 , i.e., by suitably connecting the two components on the *Comp_CG*.

In this framework, components may even be understood as independent agents, and making a handle active to a component may be understood as sending a message to the component itself. Consider the following example, representing a fragment of the code of a *controller* component/agent:

```
circuit_ok ← not fault
fault ← not fault, not test_ok
```

where *test_ok* is an incoming handle, coming from a *tester* component/agent. As soon as the *tester* will achieve *test_ok*, this incoming handle will become active, thus making the *controller* consistent, and able to conclude *circuit_ok*.

A formal definition of the methodology that we have outlined, and a detailed study of the applications, are important future directions of this research.

References

- APT, K. AND BEZEM, M., 1991. Acyclic programs. *New Generation Computing* 9 (3-4), 335–365.
- BALDUCCINI, M., BRIGNOLI, G., LANZARONE, G. A., MAGNI, F. AND PROVETTI, A., 2000. Experiments in answer sets planning. In *Proc. of the 2000 Mexican International Conference on Artificial Intelligence*.
- BALDUCCINI, M. AND GELFOND, M., 2003. Diagnostic reasoning with A-Prolog. *Theory and Practice of Logic Programming* 3(4-5), 425–461.
- BRIGNOLI, G., COSTANTINI, S., D’ANTONA, O. AND PROVETTI, A., 1999. Characterizing and computing stable models of logic programs: the non-stratified case. In *Proceedings of CIT99 Conference on Information Technology*.
- COSTANTINI, S., 1995. Contributions to the stable model semantics of logic programs with negation. *Theoretical Computer Science* 149(2), 231–255.
- COSTANTINI, S., LANZARONE, G. A., AND MAGLIOCCO, G., 1996. Asserting lemmas in the stable model semantics. In *Logic Programming: Proc. of the 1996 Joint International Conference and Symposium*, The MIT Press, Cambridge, MA, 438–452.
- COSTANTINI, S., 2001. Comparing different graph representations of logic programs under the

- answer set semantics. In *Proc. AAAI Spring Symposium "Answer Set Programming: Towards Efficient and Scalable Knowledge Representation and Reasoning"*, 21–26.
- COSTANTINI, S., D'ANTONA, O. AND PROVETTI, A., 2002. On the equivalence and range of applicability of graph-based representations of logic programs. *Information Processing Letters* 84(2), 241–249.
- COSTANTINI, S., INTRIGILA, B. AND PROVETTI, A., 2003. Coherence of updates in answer set programming. In *Proc. of the IJCAI-2003 Workshop on Nonmonotonic Reasoning, Action and Change, NRAC03*, 66–72.
- COSTANTINI, S. AND PROVETTI, A., 2004. Normal forms for answer set programming. To appear in: *Theory and Practice of Logic Programming*.
- DIX, J., 1995. A classification theory of semantics of normal logic programs: I. Strong properties. *Fundamenta Informaticae XXII*(3), 227–255.
- DIX, J., 1995. A Classification theory of semantics of normal logic programs: I. Weak properties. *Fundamenta Informaticae XXII*(3), 257–288.
- DIMOPOULOS, Y., NEBEL, B. AND KOEHLER, J., 1997. Encoding planning problems in nonmonotonic logic programs. In *Proc. of 1997 European Conference on Planning*, 169–181.
- DIMOPOULOS, Y. AND TORRES, A., 1996. Graph theoretical structures in logic programs and default theories. *Theoretical Computer Science* 170(1-2), 209–244.
- ERDEM, E. AND LIFSCHITZ, V., 2003. Tight logic programs. *Theory and Practice of Logic Programming* 3(4-5), 499–518.
- ERDOGAN, S. AND LIFSCHITZ, V., 2004. Definitions in answer set programming. In *Logic Programming and Nonmonotonic Reasoning: Proc. of 7th International Conference, LPNMR 2004*, LNAI 2923, Springer-Verlag, 114–126.
- LIFSCHITZ, V. AND TURNER, H., 1994. Splitting a logic program. In *Proceedings of the Eleventh International Conference on Logic Programming*, The MIT Press, Cambridge, MA, 23–37.
- FABER W., LEONE N. AND PFEIFER G., 1999. Pushing goal Derivation in DLP computations. In *Logic Programming and Nonmonotonic Reasoning: Proc. of 5th International Conference, LPNMR'99*, LNAI 1730, Springer-Verlag, 117–191.
- FAGES, F., 1994. Consistency of Clark's completion and existence of stable models. *Methods of Logic in Computer Science* 2, 51–60.
- FITTING, M., 1985. A Kreepe-Kleene semantics for logic programs, In: *Journal of Logic Programming* 2(4), 295–312.
- GELFOND, M. AND LIFSCHITZ, V., 1988. The stable model semantics for logic programming. In *Proceedings of the Fifth Joint International Conference and Symposium*. The MIT Press, Cambridge, MA, 1070–1080.
- GELFOND, M. AND LIFSCHITZ, V., 1991. Classical negation in logic programs and disjunctive databases. *New Generation Computing* 9, 365–385.
- KONCZAK, K., SCHAUB, T., AND LINKE, T., 2003. Graphs and colorings for answer set programming: Abridged report. In: *Answer Set Programming: Advances in Theory and Implementation, ASP03*. Volume 78 of The CEUR Workshop Proceedings Series, 137–150. <http://eur-ws.org/Vol-78/>.
- KONCZAK, K., SCHAUB, T., AND LINKE, T., 2003. Graphs and colorings for answer set programming with preferences: Preliminary report. In *Answer Set Programming: Advances in Theory and Implementation, ASP03*. Volume 78 of The CEUR Workshop Proceedings Series, 43–56. <http://eur-ws.org/Vol-78/>.
- LIBERATORE, P., 1999. Algorithms and experiments on finding minimal models. *Technical Report 09-99*, University of Rome "La Sapienza".
- LIFSCHITZ, V., 1999. Answer set planning. In *Proc. of the 1999 International Conference on Logic Programming (invited talk)*. The MIT Press, Cambridge, MA, 23–37.

- LINKE, T., 2001. Graph theoretical characterization and computation of answer sets. In: *Proceedings of IJCAI 2001*, 641–648.
- LINKE, T., 2003. Using nested logic programs for answer set programming. In *Answer Set Programming: Advances in Theory and Implementation, ASP03*. Volume 78 of The CEUR Workshop Proceedings Series, 181–194. <http://eur-ws.org/Vol-78/>.
- LINKE, T., 2003. Suitable graphs for answer set programming. In *Answer Set Programming: Advances in Theory and Implementation, ASP03*. Volume 78 of The CEUR Workshop Proceedings Series, 15–28. <http://eur-ws.org/Vol-78/>.
- MAREK, W., AND TRUSZCZYŃSKI, M., 1999. Stable models and an alternative logic programming paradigm. In *The Logic Programming Paradigm: a 25-Year Perspective*. Springer Verlag, 375–398.
- NIEMELÄ, I. 1999. Logic programs with stable model semantics as a constraint programming paradigm. *Annals of Mathematics and Artificial Intelligence* 25(3,4), 241–273, (An earlier version appeared in the Proceedings of the NM'98 workshop).
- PRZYMUSINSKA, H. AND PRZYMUSINSKI, T. C., 1990. Semantic issues in deductive databases and logic programs. R.B. Banerji (ed.) *Formal Techniques in Artificial Intelligence, a Sourcebook*, Elsevier Sc. Publ. B.V. (North Holland), 321–367.
- TURNER, H., 1994. Signed logic programs. In *Proc. of the 1994 International Symposium on Logic Programming*, 61–75.
- VAN GELDER A., ROSS K.A. AND SCHLIPF J., 1990. The well-founded semantics for general logic programs. *Journal of the ACM* 38(3), 620–650.
- WEB LOCATION OF THE MOST KNOWN ASP SOLVERS.
Cmodels: <http://www.cs.utexas.edu/users/yuliya/>
Aspps: <http://www.cs.uky.edu/ai/aspps/>
DLV: <http://www.dbai.tuwien.ac.at/proj/dlv/>
NoMoRe: <http://www.cs.uni-potsdam.de/~linke/nomore/>
Smodels: <http://www.tcs.hut.fi/Software/smodels/>